



ISSN: 0975-833X

Available online at <http://www.journalcra.com>

International Journal of Current Research  
Vol. 13, Issue, 01, pp.15548-15551, January, 2021

DOI: <https://doi.org/10.24941/ijcr.40556.01.2021>

INTERNATIONAL JOURNAL  
OF CURRENT RESEARCH

## CASE REPORT

# FUNCTIONAL AND STRUCTURAL EFFECTS OF VITAMIN D ON DIABETIC KIDNEY

**\*Burlaka Ie, A., Shevchenko T.A.**

Bogomolets National Medical University, Department of Pediatrics №4, Tolstogo str. 10, Kyiv, Ukraine, 01004

### ARTICLE INFO

#### Article History:

Received 18<sup>th</sup> October, 2020  
Received in revised form  
29<sup>th</sup> November, 2020  
Accepted 20<sup>th</sup> December, 2020  
Published online 30<sup>th</sup> January, 2021

#### Key Words:

Vitamin D, Kidney, Damage,  
T1D, Diabetic nephropathy.

### ABSTRACT

Vitamin D is vitamin and hormone with huge range of calcium-dependent and calcium-non-dependent functions. Chronic kidney disease and diabetic nephropathy patients have a high incidence of cardiovascular and infectious morbidities. Increasing evidence indicates a relationship between vitamin D deficiency and cardiovascular and infectious mortality risks. Kidney is a target organ for Vitamin D3. This article is to review and summarize the pleiotropic effects of Vitamin D in patients with Diabetes Mellitus and Diabetic Nephropathy.

Copyright © 2021, Burlaka Ie, A., Shevchenko T.A.. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Burlaka Ie, A., Shevchenko T.A. "Functional and structural effects of vitamin d on diabetic kidney", *International Journal of Current Research*, 13, (01), 15548-15551.

## INTRODUCTION

Vitamin D is a lipid-soluble vitamin and the only vitamin that can be synthesized by humans. Evolutionally, vitamin D has been synthesized by a photochemical process in land vertebrates to satisfy the requirement for a calcified skeleton for more than 350 million years [1]. Vitamin D is metabolized by 25-hydroxylase and 1 $\alpha$ -hydroxylase in the liver and kidneys, respectively, and converted to the active form, 1,25-dihydroxyvitamin D [1,25(OH)<sub>2</sub>D] [2]. Vitamin D acts through activation of the vitamin D receptor (VDR), which involves several pleiotropic effects. Level of Vitamin D synthesis and its bioavailability depends on kidney cells condition. Vitamin D deficiency is a prominent feature of kidney disease including Chronic Kidney Disease (CKD) and diabetic nephropathy (DN). Vitamin D deficiency is related to albuminuria, CKD progression, and subsequent cardiovascular diseases [3]. VDR is highly expressed in the kidney; therefore, the kidney can be considered a classic Vitamin D target organ [4]. Diabetes mellitus (DM) is a major public health problem worldwide, with ever-increasing incidence and prevalence in recent years. The Institute for Alternative Futures (IAF) expects that the total number of people with type 1 and type 2 DM in the United States will increase by 54%, from 19,629,000 to 54,913,000 people, between 2015 and 2030 [5]. Diabetic Nephropathy (DN) affects about one-third of patients with DM and currently ranks as the first cause of end-stage kidney disease in the Western world.

The major clinical manifestations of DN are proteinuria, hypertension and progressive reduction of renal function: United States Renal Data System registered that approximately 35–50% of end-stage renal disease (ESRD) cases are secondary to DN complications [6]. Thus, studies of molecular mechanisms preserving kidneys which Vitamin D involved in are important in terms of prevention of proteinuric kidney diseases, i.e. Diabetic Nephropathy.

**Functional effects of Vitamin D on diabetic kidney:** The complexity of interactions of Vitamin D is directly related with progressive long-term changes implicated in the worsening of renal function. These changes result in a dysregulation of the Vitamin D-dependent pathways. Various studies demonstrated a pivotal role of Vitamin D supplementation in regression of albuminuria and glomerulosclerosis, contrasting the increase of glomerular basement membrane thickening and podocyte effacement. These, in turn, improves renal and cardiovascular outcomes. The homeostasis and regulation of the nephron's function are absolutely dependent from the cross-talk between endothelium and podocytes. A pivotal role for the management of DM could be found in the pleiotropic actions of vitamin D. Studies conducted by Afzal et al. [7], Schöttker et al. [8], Tsur et al. [9], have shown a link between low serum levels of vitamin D and the increasing risk of developing DM. An association of a proper status of vitamin D with a better modulation of glucose homeostasis, due to a regulation of insulin secretion and tyrosine phosphorylation of the insulin receptor shown [10]. Data of the Third National Health and Nutrition Examination

\*Corresponding author: Burlaka Ie, A.,

Bogomolets National Medical University, Department of Pediatrics №4, Tolstogo str. 10, Kyiv, Ukraine, 01004

Survey (NHANES III) found an inverse association between the level of vitamin D and the prevalence of albuminuria. In our previous studies conducted in Department of Pediatrics №4 of Bogomolets National Medical University we found decreased level of Vitamin D in iabetic patients. In group with T1D Vitamin D level was in insufficiency rage, in patients with DN Vitamin D deficiency found [11,12].

The study of Karnchanasorn et al. [13] showed modulatory effect of vitamin D. Authors demonstrated a correlation between appropriate vitamin D status and both  $\beta$ -cell function and insulin sensitivity. A protective role of vitamin D in type 1 DM has been demonstrated in a recent where serum 25-(OH) D concentrations sustained with cholecalciferol supplementation 3000 IU/day for one year improved metabolic control and slowed the decline of residual  $\beta$ -cell function in children with T1DM [14]. The meta-analysis by Gregoriou et al. [15] also showed that vitamin D supplementation in the form of 1 $\alpha$ (OH)vitamin and cholecalciferol appears to be beneficial in the treatment of T1DM patients by attenuating the natural history of the disease. Activation of the VDR is essential in reducing proteinuria. Traditionally, administration of RAAS blockers can reduce albuminuria [16]. 1.25(OH)<sub>2</sub>D<sub>3</sub> is known as a RAS inhibitor by its negative regulatory effect on renin production to provide additional renoprotection [17].

**Structural effects of Vitamin D on diabetic kidney:** The renoprotective effects of vitamin D can improve proteinuria, glomerulosclerosis, and interstitial infiltration and reduce renal oxidative stress [8]. Combined treatment with paricalcitol and losartan suppressed the induction of fibronectin, transforming growth factor  $\beta$  (TGF- $\beta$ ) and monocyte chemoattractant protein-1 (MCP-1), and reversed the decline of the slit diaphragm proteins nephrin, Neph-1, ZO-1, and alpha-actinin-4 [19]. Another emerging issue linked with DN development and lower levels of Vitamin D is urinary megalin excretion in T2DM patients. [19]. Megalin, as a member of the low-density lipoprotein receptor family [20]. Megalin plays an important role in the reabsorption of Vitamin D binding protein from glomerular filtrates. It was reported that megalin-mediated (auto)lysosomal dysfunction in primary tubular epithelial cells is decisive for the development of kidney disease in a High-Fat-Diet-induced diabetes model. Exocytosis-mediated urinary megalin excretion increases along with the progression of DN, giving further contributions in the understanding of the pathogenesis of Vitamin D loss in these subjects, their findings also suggest a potential role of megalin urinary excretion as indicator of progression of DN [22,23].


CKD and DN always associated with high rate of cardiovascular complications. There are data showing that paricalcitol and calcitriol are both able to modulate the thrombomodulin expression in human aortic smooth muscle cells. Moreover, D'Arrigo et al. demonstrated a raise of soluble thrombomodulin levels in patients with CKD by the administration of paricalcitol. [24,25]. VDR knockout in diabetic mice was associated with severe albuminuria and glomerulosclerosis. Alternatively, vitamin D might slow the progression of diabetic nephropathy by improving insulin secretion, delaying destruction of  $\beta$  islet cells, affecting osteocalcin, and consequently assisting in glucose metabolism. TGF- $\beta$ , MCP-1, hepatocyte growth factor, thrombospondin-1, and plasminogen activator inhibitor are other possible molecular targets of vitamin D action [26]. Increasing

evidence from experimental and clinical studies has unveiled a pathological role of macrophages in the development of glomerulosclerosis by the production of inflammatory chemokines, cytokines and fibrogenic factors, release of proteolytic enzymes and production of reactive oxygen species. A central role in these mechanisms is played by the monocyte chemoattractant protein (MCP)-1, a chemokine produced by mesangial cells (MCs) and renal tubular cells that has the responsibility of recruiting macrophages into the kidney [27]. It was demonstrated that calcitriol can inhibit the synthesis and activity of MCP-1 and contrast the glomerular injury in diabetic mice. Their data suggest that vitamin D may protect against renal injury in DN by preventing or reducing macrophage infiltration [27,28]. A proper vitamin D regulation also has an important role against inflammation secondary to DN. Indeed, diabetes leads to an increase in the expression of inflammatory factors and inappropriate immune activity. Inflammatory response likely contributes to DM occurrence by worsening insulin resistance and it is in turn intensified in the presence of hyperglycaemia to exacerbate long-term complications of diabetes [29].

Vitamin D exerts protective effects against inflammatory agents by inhibiting the expression of interleukin (IL)-6, IL-8, RANTES (regulated on activation, normal T cell expressed and secreted), intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), platelet-endothelial cell adhesion molecule-1 (PECAM-1), receptor of advanced glycation end products (RAGE) and E-selectin through a nuclear factor- $\kappa$ B (NF- $\kappa$ B)-mediated mechanism, partly by disrupting DNA binding of NF- $\kappa$ B. Moreover, vitamin D represses the expression of cyclooxygenase (COX)-2 and upregulates the expression of 15-hydroxyprostaglandin dehydrogenase (15PGDH), the enzyme initiating prostaglandin catabolism, in this way reducing prostaglandin levels and suppressing the production of several proinflammatory cytokines. The immunomodulatory action of vitamin D has been well demonstrated also in the study of Lucisano et al., where an acute paricalcitol supplementation induced a significant reduction of IL-17, IL-6, IL-1 $\beta$ , TNF- $\alpha$  and IFN- $\gamma$  in a cohort of CKD patients [30]. Chronic inflammation leads to sclerotic processes in kidney. Reduction of the inflammatory rate can prevent irreversible morphological changes in kidneys. In the study conducted on human incubated monocytes taken from patients with DM and DN with uraemia, it was found that Vitamin D may exert an anti-inflammatory effect by regulating the signal transduction pathways that control VDR and signal transducer and activator of transcription 5 (STAT5) expression.

These findings have been further investigated. It was found more anti-inflammatory effects of Vitamin D and VDR on phosphorylated STAT5 (p-STAT5) in serum-incubated monocytes from patients with DM and uraemia caused by DN: lipopolysaccharide associated with IL-15 upregulated the expression of p-STAT5, whereas pre-treatment with 1,25-(OH)<sub>2</sub>D<sub>3</sub> significantly inhibited this effect [31]. Experimental as well as observational studies and clinical trials conducted in the past years suggest the effective role of Vitamin D and the synergic action with RAAS inhibitors to counteract the worsening of DN and to preserve the glomeruli and the integrity of glomerular filtration barrier. Moreover, Vitamin D seems to exert many extra-renal functions essential for the body homeostasis. The results of these studies emphasize the need for better awareness among researchers and clinicians

about the consequences of insufficient Vitamin D levels and the importance of monitoring its status in high-risk populations. Summarized scheme of Vitamin D effects in Diabetes Mellitus and Diabetic nephropathy given in Picture 1.



**Picture 1. Summarized scheme of Vitamin D in Diabetes Mellitus and Diabetic nephropathy**

Even if growing evidence proves that vitamin D may have antiproteinuric, anti-inflammatory, renoprotective and cardioprotective effects in patients with DN, it is still required randomized controlled trials in larger patient groups studying molecular background of the nephroprotective effects of Vitamin D.

## REFERENCES

- Afzal S., Bojesen S.E., Nordestgaard, B.G. Low 25-hydroxyvitamin D and risk of type 2 diabetes: A prospective cohort study and metaanalysis. *Clin. Chem.* – 2013. - 59:381–391.
- Alicic R.Z., Rooney M.T., Tuttle K.R. Diabetic Kidney Disease: Challenges, Progress, and Possibilities. *Clin J Am Soc Nephrol.* – 2017. - 12(12):2032-2045.
- Anavekar N.S., Gans D.J., Berl T., Rohde R.D., Cooper W., Bhaumik A., Hunsicker L.G., Rouleau J.L., Lewis J.B., Rosendorff C., et al. Predictors of cardiovascular events in patients with type 2 diabetic nephropathy and hypertension: A case for albuminuria. *Kidney Int. Suppl.* – 2004. - 50–55.
- Bikle D. Vitamin D: Production, Metabolism, and Mechanisms of Action. [Updated 2017 Aug 11]. In: Feingold KR, Anawalt B, Boyce A, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK278935/>
- Brenner B.M., Cooper M.E., de Zeeuw D., Keane W.F., Mitch W.E., Parving H.H., Remuzzi G., Snapinn S.M., Zhang Z., Shahinfar S. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N. Engl. J. Med.* – 2001. - 345:861–869.
- Burlakale. Beyond Proteinuria: Apoptosis and Vitamin D3 State in Children with Diabetic Nephropathy. *J Clin Exp Pathol.* – 2016. - 6: 266.
- D'arrigo G., Pizzini P., Cutrupi S., Tripepi R., Tripepi G., Mallamaci F., Zoccali C. Vitamin D receptor activation raises soluble thrombomodulin levels in chronic kidney disease patients: A double blind, randomized trial. *Nephrol. Dial. Transplant.* – 2018. -doi: 10.1093/ndt/gfy104.FP348.
- De S., Kuwahara S., Hosojima M., Ishikawa T., Kaseda R., Sarkar P., Yoshioka Y., Kabasawa H., Iida T., Goto S., et al. Exocytosis-Mediated Urinary Full-Length Megalin Excretion Is Linked with the Pathogenesis of Diabetic Nephropathy. *Diabetes.* – 2017. - 66:1391–1404.
- De Zeeuw D., Remuzzi G., Parving H.H., Keane W.F., Zhang Z., Shahinfar S., Snapinn S., Cooper M.E., Mitch W.E., Brenner B.M. Proteinuria, a target for renoprotection in patients with type 2 diabetic nephropathy: Lessons from RENAAL. *Kidney Int.* – 2004. - 65:2309–2320.
- DeLuca H.F. Overview of general physiologic features and functions of vitamin D. *Am J Clin Nutr.* – 2004. - 80(6 Suppl): 1689-1696.
- Diez-Sampedro A., Lenz O., Forni A. Podocytopathy in diabetes: A metabolic and endocrine disorder. *Am. J. Kidney Dis.* – 2011. - 58:637–646.
- Flores M. A role of vitamin D in low-intensity chronic inflammation and insulin resistance in type 2 diabetes mellitus? *Nutr. Res. Rev.* – 2005. - 18:175–182.
- Franca Gois P.H., Wolley M., Ranganathan D., Seguro A.C. Vitamin D Deficiency in Chronic Kidney Disease: Recent Evidence and Controversies. *Int J Environ Res Public Health.* – 2018. -15(8):1773.
- Gembillo G., Cernaro V., Salvo A., et al. Role of Vitamin D Status in Diabetic Patients with Renal Disease. *Medicina (Kaunas).* – 2019. - 55(6):273.
- Gregoriou E., Mamais I., Tzanetakou I., Lavranos G., Chrysostomou S. The Effects of Vitamin D Supplementation in Newly Diagnosed Type 1 Diabetes Patients: Systematic Review of Randomized Controlled Trials. *Rev. Diabet. Stud.* – 2017. - 14:260–268.
- Hammer Y., Soudry A., Levi A., Talmor-Barkan Y., Leshem-Lev D., Singer J., Kornowski R., Lev E.I. Effect of vitamin D on endothelial progenitor cells function. *PLoS ONE.* – 2017. – 12.
- Henry H.L. Regulation of vitamin D metabolism. *Best Pract Res Clin Endocrinol Metab.* – 2011. - 25(4):531-541.
- Karnchanasorn R., Ou H.Y., Chiu K.C. Plasma 25-hydroxyvitamin D levels are favorably associated with  $\beta$ -cell function. *Pancreas.* – 2012. - 41:863–868.
- Lontchi-Yimagou E., Sobngwi E., Matsha T.E., Kengne A.P. Diabetes mellitus and inflammation. *Curr. Diab. Rep.* – 2013. - 13:435–444.
- Lucisano S., Arena A., Stassi G., Iannello D., Montalto G., Romeo A., Costantino G., Lupica R., Cernaro V., Santoro D., Buemi M. Role of Paricalcitol in Modulating the Immune Response in Patients with Renal Disease. *Int. J. Endocrinol.* – 2015. - 2015:765364.
- Maestroni S., Zerbini G. Glomerular endothelial cells versus podocytes as the cellular target in diabetic nephropathy. *Acta Diabetol.* - 2018 doi: 10.1007/s00592-018-1211-2.
- Maidannyk V., Burlakale. Association of Vitamin D deficiency, cellular hypoxia, and caspase-3 with renal disease in pediatric diabetic nephropathy. *Journal of Translational Science.* – 2016. -2(2): 130-133.
- Ogasawara S., Hosojima M., Kaseda R., Kabasawa H., Yamamoto-Kabasawa K., Kurosawa H., Narita I. Significance of urinary full-length and ectodomain forms of megalin in patients with type 2 diabetes. *Diabetes Care.* – 2012. - 35:1112–1118.
- Panjiyar R.P., Dayal D., Attri S.V., Sachdeva N., Sharma R., Bhalla A.K. Sustained serum 25-hydroxyvitamin D concentrations for one year with cholecalciferol

- supplementation improves glycaemic control and slows the decline of residual  $\beta$  cell function in children with type 1 diabetes. *Pediatr. Endocrinol. Diabetes Metab.* – 2018. - 111–117.
- Pilz S., van Den Hurk K., Nijpels G., Stehouwer C.D.A., Van'triet E., Kienreich K., Dekker J.M. Vitamin D status, incident diabetes and prospective changes in glucose metabolism in older subjects: The Hoorn study. *Nutr. Metab. Cardiovasc. Dis.* – 2012. - 22:883–889.
- Saito A., Pietromonaco S., Loo A.K., Farquhar M.G. Complete cloning and sequencing of rat gp330/‘megalin’ a distinctive member of the low density lipoprotein receptor gene family. *Proc. Natl. Acad. Sci. USA.* – 1994. - 91:9725–9729.
- Schöttker B., Herder C., Rothenbacher D., Perna L., Müller H., Brenner H. Brenner Serum 25-hydroxyvitamin D levels and incident diabetes mellitus type 2: A competing risk analysis in a large population-based cohort of older adults. *Eur. J. Epidemiol.* – 2013. - 28:267–275.
- Tsur B.S., Feldman I., Feldhammer M.B., Hoshen G., Leibowitz R.D. Balicer Decreased serum concentrations of 25-hydroxycholecalciferol are associated with increased risk of progression to impaired fasting glucose and diabetes. *Diabetes Care.* – 2013. - 36:1361–1367.
- Yang M., Yang B.O., Gan H., Li X., Xu J., Yu J., Li F. Anti-inflammatory effect of 1,25-dihydroxyvitamin D3 is associated with crosstalk between signal transducer and activator of transcription 5 and the vitamin D receptor in human monocytes. *Exp. Ther. Med.* – 2015. - 9:1739–1744.
- Zhang Z, Sun L, Wang Y, Ning G, Minto AW, Kong J, Quigg RJ, Li YC. Renoprotective role of the vitamin D receptor in diabetic nephropathy. *Kidney Int.* – 2008. - 73(2):163–171.
- Zhang Z., Yuan W., Sun L., Szeto F.L., Wong K.E., Li X., Kong J., Li Y.C. 1,25-Dihydroxyvitamin D3 targeting of NF- $\kappa$ B suppresses high glucose-induced MCP-1 expression in mesangial cells. *Kidney Int.* – 2007. - 72:193–201.

\*\*\*\*\*