




# Distribution, preparedness and management of Ukrainian adult refugees on dialysis—an international survey by the Renal Disaster Relief Task Force of the European Renal Association

Ewa Pawłowicz-Szlarska<sup>1</sup>, Raymond Vanholder<sup>2,3</sup>, Mehmet S. Sever<sup>4</sup>, Serhan Tuğlular<sup>5</sup>, Valerie Luyckx<sup>6,7,8</sup>, Kai-Uwe Eckardt<sup>9</sup>, Daniel Gallego<sup>3,10</sup>, Dmytro Ivanov<sup>11</sup>, Ionut Nistor<sup>12,13</sup>, Rukshana Shroff<sup>14</sup>, Andrej Škoberne<sup>15,16</sup>, Stefano Stuard<sup>17</sup>, Ryszard Gellert<sup>18</sup>, Edita Noruišiene<sup>3,19</sup>, Mohamed Sekkarie<sup>20</sup> and Andrzej Wiecek<sup>21</sup>; on behalf of the Renal Disaster Relief Task Force of the ERA

<sup>1</sup>Department of Nephrology, Hypertension and Kidney Transplantation, Medical University of Łódź, Łódź, Poland, <sup>2</sup>Nephrology Section, Department of Internal Medicine and Pediatrics, Ghent University Hospital, Ghent, Belgium, <sup>3</sup>European Kidney Health Alliance, Brussels, Belgium, <sup>4</sup>Department of Nephrology, Istanbul School of Medicine, Istanbul University, Istanbul, Turkey, <sup>5</sup>Department of Nephrology, School of Medicine, Marmara University, Istanbul, Turkey, <sup>6</sup>Renal Division, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA, <sup>7</sup>Department of Paediatrics and Child Health, University of Cape Town, Cape Africa, South Africa, <sup>8</sup>Department of Public and Global Health, Epidemiology, Biostatistics and Prevention Institute, University of Zurich, Zurich, Switzerland, <sup>9</sup>Department of Nephrology and Medical Intensive Care, Charité-Universitätsmedizin Berlin, Berlin, Germany, <sup>10</sup>European Kidney Patient Federation, Wien, Austria, <sup>11</sup>Department of Nephrology and RRT, Bogomolets National Medical University, Kyiv, Ukraine, <sup>12</sup>Department of Internal Medicine, Nephrology and Geriatrics, Grigore T. Popa University of Medicine and Pharmacy, Iasi, Romania, <sup>13</sup>Department of Nephrology, Dr C. I. Parhon University Hospital, Iasi, Romania, <sup>14</sup>Renal Unit, UCL Great Ormond Street Hospital and Institute of Child Health, London, UK, <sup>15</sup>Department of Nephrology, Division of Internal Medicine, University Medical Centre Ljubljana, Ljubljana, Slovenia, <sup>16</sup>Medical Faculty, University of Ljubljana, Ljubljana, Slovenia, <sup>17</sup>Global Medical Office, Fresenius Medical Care, Bad Homburg, Germany, <sup>18</sup>Department of Nephrology and Internal Medicine, Centre for Postgraduate Medical Education, Warsaw, Poland, <sup>19</sup>European Dialysis and Transplant Nurses Association – European Renal Care Association, Hergiswil, Switzerland, <sup>20</sup>Nephrology and Hypertension Associates, Bluefield, WV, USA and <sup>21</sup>Department of Nephrology, Transplantation and Internal Medicine, Medical University of Silesia, Katowice, Poland

Correspondence to: Andrzej Wiecek; E-mail: [awiecek@sum.edu.pl](mailto:awiecek@sum.edu.pl)



**ABSTRACT**

**Background.** Due to the Russian–Ukrainian war, some of the about 10 000 adults requiring dialysis in Ukraine fled their country to continue dialysis abroad. To better understand the needs of conflict-affected dialysis patients, the Renal Disaster Relief Task Force of the European Renal Association conducted a survey on distribution, preparedness and management of adults requiring dialysis who were displaced due to the war.

**Methods.** A cross-sectional online survey was sent via National Nephrology Societies across Europe and disseminated to their dialysis centers. Fresenius Medical Care shared a set of aggregated data.

**Results.** Data were received on 602 patients dialyzed in 24 countries. Most patients were dialyzed in Poland (45.0%), followed by Slovakia (18.1%), Czech Republic (7.8%) and Romania (6.3%). The interval between last dialysis and the first in the reporting center was 3.1 ± 1.6 days, but was ≥4 days in 28.1% of patients. Mean age was 48.1 ± 13.4 years, 43.5% were females. Medical records were carried by 63.9% of patients, 63.3% carried a list of medications, 60.4% carried the medications themselves and 44.0% carried their dialysis prescription, with 26.1% carrying all of these items and 16.1% carrying none. Upon presentation outside Ukraine, 33.9% of patients needed hospitalization. Dialysis therapy was not continued in the reporting center by 28.2% of patients until the end of the observation period.

**Conclusions.** We received information about approximately 6% of Ukrainian dialysis patients, who had fled their country

by the end of August 2022. A substantial proportion were temporarily underdialyzed, carried incomplete medical information and needed hospitalization. The results of our survey may help to inform policies and targeted interventions to respond to the special needs of this vulnerable population during wars and other disasters in the future.

**Keywords:** conflicts, dialysis, disasters, refugees, wars

**INTRODUCTION**

On 24 February 2022 Russia initiated an invasion of Ukraine [1]. As of 24 September 2022, the number of Ukrainians who fled the country was estimated to be almost 7.5 million [2]. Additionally, several millions of Ukrainians migrated internally, mostly to the western part of Ukraine [3].

The impact of conflicts on health is overwhelming [4]. Direct effects include gunshot injuries and trauma-related morbidity and mortality. Assessment of indirect effects is challenging, as they remain underestimated and under-recorded. They include the impact of conflict on infectious diseases, non-communicable diseases, morbidity due to reduced healthcare access, and interruption of antenatal, pediatric and maternal services and of mental healthcare. In recent years, there has been a growing awareness of the burden of chronic diseases in conflicts [4].

There are several reports on the situation of patients with kidney diseases during natural disasters [5, 6], but less is known about the influence of military conflicts on this vulnerable

## KEY LEARNING POINTS

### What is already known about this subject?

- During armed conflicts dialysis patients may experience limitations or interruptions to therapy leading to severe life-threatening complications due to medical and logistical challenges.
- Before the Russian–Ukrainian war, there were approximately 10 000 adults receiving long-term dialysis in Ukraine.
- Some of these patients fled their place of residence and looked for opportunities to continue dialysis elsewhere in Ukraine or abroad. The number of such patients and their condition after displacement were thus far unknown.

### What this study adds?

- By the end of August 2022, we received information about approximately 6% of Ukrainian dialysis patients who fled their country since the start of the Russian–Ukrainian conflict, with the majority receiving dialysis in a country neighboring Ukraine.
- Preparedness for displacement varied and was incomplete in most patients.
- Almost 30% of displaced patients remained without dialysis for 4 days or longer.
- Clinical status upon arrival to the host countries often necessitated hospitalization.

### What impact this may have on practice or policy?

- Monitoring the movement and condition of dialysis patients during armed conflicts and the present results may help inform policies that allow for prompt and targeted interventions to respond to the special needs of this vulnerable population during wars and other emergencies around the globe.

population. The impact of wars on patients with kidney diseases was recently summarized in a consensus statement from the Renal Disaster Relief Task Force (RDRTF) of the European Renal Association (ERA) [7]. Patients on maintenance hemodialysis (HD) are at risk of prolonged underdialysis due to damaged healthcare infrastructure, suboptimal availability of water and energy supplies, shortage of dialysis personnel, transportation difficulties or the need to flee. Consequently, especially non-adherence to fluid and dietary restrictions may lead to serious, life-threatening complications. Patients on peritoneal dialysis (PD) may face logistical problems in obtaining dialysis solutions on time, and are at high risk of complications, such as peritoneal dialysis-related peritonitis or exit-site infections due to unhygienic circumstances. Patients after kidney transplantation may suffer from interruptions of maintenance immunosuppressive treatment and are at risk of serious infections due to their immunosuppressed state [7].

The possibility of leaving affected zones may be impacted by many factors including severity of disability, and economic, personal, emotional and cultural determinants [7].

As of 1 January 2021, there were 11 181 patients (268 per million population) on kidney replacement therapy (KRT) in Ukraine, including 6017 on HD, 2700 on hemodiafiltration, 931 on PD and 1533 with kidney transplants [8]. The exact number of Ukrainian KRT patients who were internally displaced, mostly to western parts of Ukraine, or who fled to neighboring countries is currently unknown. In accordance with the European Union protection act [9] neighboring country governments and healthcare systems provided support, including full reimbursement of healthcare costs and procedural assistance outside Ukraine. In addition, international kidney care organizations (ERA, the American Society of Nephrology, the International Society of Nephrology and the European Kidney Health Alliance) supported by the World Health Organization joined together in these relief

efforts, including supply of dialysis equipment to Ukraine, organization of fundraisers, assistance for kidney patients remaining in Ukraine and to those displaced outside the country, as well as various support actions for Ukrainian nephrologists [10].

Due to scarce epidemiological data on refugees suffering from kidney failure, the RDRTF of the ERA conducted an international survey on distribution, preparedness and management of adults requiring dialysis outside Ukraine since the beginning of the war. The results may lead to a better understanding of the needs of patients with kidney failure affected by a conflict and may support improvement in preparedness procedures and equitable and evidence-based allocation of resources for this vulnerable population in ongoing and future conflicts.

## MATERIALS AND METHODS

### Data source

#### Survey data

A web-based survey hosted on the Survey Monkey application was distributed across Europe and adjacent countries that are located in the geographic area covered by the ERA (list of the countries is provided in Supplementary data, Table S1). The first e-mail invitation was sent to Presidents of National Societies of Nephrology on 16 May 2022 along with a request to disseminate the survey to a lead physician of each dialysis unit within their countries. The invitation was repeated after 5 weeks. The survey requested data about adult ( $\geq 18$  years old) dialysis patients displaced from Ukraine, who received dialysis in public or private dialysis centers. Data collection was anonymized making identification of individual patients impossible. Details of the survey are provided in Supplementary data, File S1.

### Aggregated data

Due to internal privacy procedures, all dialysis units from Fresenius Medical Care (FMC) shared a set of aggregated data without direct center participation. Results from all FMC European, the Middle East and Africa (EMEA) region units were provided by the FMC representatives.

### Data collection

Data were collected between May 2022 and August 2022. The survey comprised questions on dialysis therapy before and at the time of war both in Ukraine and in the other countries, patients' preparedness for displacement outside Ukraine, hospitalizations outside Ukraine, coronavirus disease 2019 (COVID-19) history, language barrier-related issues and psychological support received by the patients, demographic features (age, sex, education level) and medical history (cause of kidney failure). Questions about dialysis therapy concerned dialysis modality, time since start of dialysis, vascular access type, viral status and pharmacological treatment (erythropoiesis-stimulating agents, intravenous iron, vitamin D analogues, cinacalcet and phosphate binders).

Possible interruptions of treatment in Ukraine since the start of war (i.e. decreased frequency of dialysis sessions) were also addressed. Data about countries where patients were dialyzed after leaving their regular units in Ukraine were reported as well as data about kidney transplantation waitlist status in Ukraine and in the host country.

Additionally, for the cohort of patients dialyzed in Poland, the country hosting most Ukrainian dialysis patients, clinical and laboratory data at the time of admission to the center were gathered. Blood pressure and interdialytic body weight gain on admission to the reporting center as well as the first reported laboratory parameters were collected. Data about war-related or journey-related injuries were also gathered.

Aggregated data sent by FMC comprised gender, mean age, duration of KRT, viral status, vascular access type, COVID-19 testing and pharmacotherapy at the receiving center. The survey response rate for Polish dialysis units was 85%. For FMC EMEA units it was 100% (related to the aggregated nature of the received data).

The survey database was checked for double entries but no such cases were found.

All percentages sum up to 100%, although the total number of responses included in the percentage calculations was different for each question due to different number of non-responses and because some questions were covered only by the point-by-point survey and not by the FMC data. For some questions, 'No data' was an answer option—for those questions these responses were included up to 100%. The list of non-responses per particular question is presented in Supplementary data, Table S2.

### Study oversight

The study was designed by the RDRTF of ERA. The study protocol and the survey were reviewed and approved by the ERA Scientific Advisory Board. The questionnaire was

**Table 1: Number of dialysis patients displaced outside Ukraine and dialyzed abroad.**

Country	Number of patients		Total
	Survey data	Aggregated data	
Poland	161	110	271
Slovakia	2	107	109
Czech Republic	19	28	47
Romania	9	29	38
Germany	28	0	28
Hungary	6	15	21
Russia	0	12	12
Switzerland	11	0	11
Lithuania	9	0	9
Georgia	5	0	5
Latvia	5	0	5
Denmark	4	0	4
Belgium	3	0	3
Israel	3	0	3
Netherlands	2	0	2
Croatia	1	0	1
Finland	1	0	1
Luxembourg	1	0	1
Slovenia	1	0	1
Italy, Estonia, France, Portugal, Spain	Italy—14  Estonia—5 France—0 Portugal—0 Spain—0	11 <sup>a</sup>	30

<sup>a</sup>Due to the internal privacy policy, FMC gave the total number of patients for these five countries.

submitted to the Ethics Committee of the Medical University of Silesia in Katowice, Poland and need for ethics review was waived. Detailed authors' contributions are provided at the end of this article.

## RESULTS

Data on 602 patients who were dialyzed abroad in 24 countries after leaving Ukraine are reported. The distribution of patients per country is provided in Table 1. Most patients were dialyzed in Poland (45.0%), followed by Slovakia (18.1%), Czech Republic (7.8%), Romania (6.3%), Germany (4.7%) and Hungary (3.5%) (survey and aggregated FMC data together).

The location of dialysis units in Ukraine where patients were dialyzed before the displacement is provided in Fig. 1. Besides Kyiv, most patients left centers located in the south-east of Ukraine (survey data only).

At least one dialysis session/visit in one or more dialysis centers between the home institution in Ukraine and the reporting center had occurred in 34.6% of patients. The largest group of patients had these transit dialysis sessions in Poland (41.3%), Ukraine (25%, mostly in Lviv) or Slovenia (7.5%). All patients treated ad interim in Slovenia were finally dialyzed in Italy (survey data only).

Mean patient age was  $48.1 \pm 13.4$  years, and 43.5% were females. Regarding education levels in the study group, 14.6% of patients had a university degree, 8.9% had vocational education, 18.9% secondary education and 3.2% primary education (data about education were captured only from



**Figure 1:** Location of dialysis units in Ukraine, where patients were dialyzed before the displacement and number of these patients.

**Table 2: Clinical data with regard to vascular access type (HD patients) and viral status (HD and PD patients) in the study group.**

	% of patients
Vascular access type	
Arteriovenous fistula	89
Arteriovenous graft	2.0
Tunneled catheter	5.1
Non-tunneled catheter	3.9
Viral status	
HBs antigen positive	7.3
Anti-HCV antibodies	16.1
Anti-HIV antibodies	0.6
Anti-HBc antibodies	27.3

HBs antigen, antigen of the hepatitis B virus; HCV, hepatitis C virus; HIV, human immunodeficiency virus; HBc antigen, hepatitis B virus protein (core antigen).

45.6% of migrants). Before leaving Ukraine, 98% had been on HD, 1.2% on continuous ambulatory PD and 0.8% on automated PD. Therapeutic shift from PD to HD after displacement occurred in two patients. Mean dialysis vintage was  $57.7 \pm 66.6$  months (mean  $\pm$  SD). Data on dialysis vascular access type (HD patients) and viral status (all patients) are provided in Table 2 (survey and aggregated data). The majority (89%) of HD patients had an arteriovenous fistula. Hepatitis B (HBs) antigen positivity was noted in 7.3% and anti-HBc antibodies in 27.3% of all patients.

Primary causes of kidney failure are presented in Fig. 2 (survey data only). The most common causes of kidney disease were glomerulonephritis and autoimmune diseases (29.5%). In 20.7% the cause was unknown.

Under war conditions in Ukraine, HD session frequency was reduced in 23.5% of those patients who finally migrated.

Mean time between last dialysis session and the first session in the reporting center was  $3.1 \pm 1.6$  days. In 28.1% of patients this time interval ranged from 4 (16.8%) up to 12 days (survey data only).


Over one-quarter (28.2%) of patients had not continued dialysis therapy in the reporting center by the closure date of the survey (31 August). Mean number of HD visits was  $11.6 \pm 11.2$ ; median 8, range 1–50; only three PD patients left the reporting centers, but the duration of stay was not provided. Of those not continuing in the reporting center, 4.1% had died and 94.6% left for another country or another city in the same country. Of the patients who left, 28.6% intended to return to Ukraine (survey data only).

Patient preparedness for displacement is outlined in Table 3. Overall, 26.1% of the patients presented in the host dialysis unit with their medical history, medication list, medications and dialysis prescriptions, whereas 16.1% presented with none of these (survey data only).

After leaving Ukraine, 33.9% of patients were hospitalized upon presentation at the reporting center. The most common causes of hospitalizations are presented in Table 4.

In the reporting dialysis units, 53.6% of patients received erythropoiesis-stimulating agents, 39.4% intravenous iron supplementation, 37.5% phosphate binders, 22.2% vitamin D analogues and 3.5% cinacalcet (survey and aggregated data). However, data on pharmacotherapy received in Ukraine before the displacement was insufficiently available to allow reliable information to be reported.

Twenty-nine patients (11.5%) had been waitlisted for kidney transplantation in Ukraine. Of these, 15 were also waitlisted for transplantation in their host country. In addition,



**Figure 2:** Primary causes of kidney failure in the study group (survey data only). ADPKD, autosomal dominant polycystic kidney disease; ARPKD, autosomal recessive polycystic kidney disease; CAKUT, congenital anomalies of the kidney and urinary tract.

**Table 3: Possession of particular determinants of preparedness for displacement with regard to medical safety (survey data only).**

Preparedness determinant	% of patients
Medical records	63.9
List of medications	63.3
Medications themselves	60.4
Dialysis prescription	44.0

**Table 4: Causes of hospitalizations outside Ukraine among displaced dialysis patients.**

Cause of hospitalization	Number of patients	% of patients
Anemia	16	18.6
COVID-19 pneumonia	4	4.7
Non-COVID-19 pneumonia	4	4.7
Fluid overload	4	4.7
Need for tunneled catheter implantation	4	4.7
Heart failure	3	3.5
Peripheral artery occlusive disease	3	3.5
Need AVF surgery	3	3.5
Acute cholecystitis	2	2.3
Above knee amputation	2	2.3
Myeloma	2	2.3
Endocarditis	2	2.3
Arterial hypertension	2	2.3
Dialysis peritonitis	2	2.3
Arrhythmia	2	2.3
AVF thrombosis	2	2.3
GI hemorrhage	2	2.3
Ileus	2	2.3

AVF, arteriovenous fistula; GI, gastrointestinal.

33 patients who had not been listed for transplantation in Ukraine started the listing process under the care of the reporting center (survey data only).

Of the 90.5% of patients tested for active severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in the reporting center, 2.1% were positive (survey and aggregated data). At the time of presentation to the reporting center, 32.1% of patients had not been vaccinated, 8% had received three doses, 29.3% two doses and 1.6% one dose. Twenty-six patients (10.6%) were vaccinated while being cared for at the reporting dialysis center (survey data only).

Fifty-four percent of patients travelled to the destination country accompanied by at least one other person, mostly family members or, rarely, other people also requiring dialysis (survey data only).

Communication and language challenges were reported by 50.6% of responding physicians. Translation was not required in 30.4% of cases, in 26.4% a coworker was able to translate, in 17.2% patient's family members helped in translation, while 16.4% of responding physicians used telephone/internet translation services. In 9.6% of cases an interpreter, provided by hospital or dialysis unit, was physically available on site (survey data only).

A total of 43 patients (17.2%) received psychological support while being cared for in the reporting dialysis center. This service was provided by the dialysis unit itself for 29 patients and outside the center for 14 patients (survey data only).

**Table 5: Laboratory parameters at admission to the hosting dialysis units collected in Poland.**

Parameter	N	Mean	SD	Median	Min-max	Percentage	
						Range	% (N)
Hemoglobin (g/dL)	126	9.5	1.8	9.15	5–14	<7.0	7.9 (10)
						7.0–7.9	7.9 (10)
						8.0–8.9	25.4 (32)
						9.0–9.9	23.8 (30)
						10.0–11.5	19 (24)
>11.5	15.9 (20)						
WBC count ( $\times 10^3/\mu\text{L}$ )	118	6.6	2.4	6.3	2.7–16.3		
Platelet count ( $\times 10^3/\mu\text{L}$ )	119	209.7	67.4	203	86–542		
Potassium (mmol/L)	120	5.4	0.8	5.3	3.0–7.9	<3.5	0.8 (1)
						3.5–5.0	30.8 (37)
						5.1–6.0	45.8 (55)
						6.1–7.0	20.8 (25)
						>7.0	1.7 (2)
Sodium (mmol/L)	113	138.7	2.7	139	132–145		
Calcium (mmol/L)	98	2.2	0.3	2.2	1.6–3.1		
Phosphorus (mmol/L)	93	3.6	2.3	2.4	0.9–10.1		
Creatinine ( $\mu\text{mol/L}$ )	74	897.7	262.2	900.8	450.8–1522.2		
BUN (mg/dL)	97	75.2	29.3	72.4	28–197		

BUN, blood urea nitrogen; SD, standard deviation; WBC, white blood cells.

### Clinical data of the Polish cohort (survey data only)

Data about volume status, blood pressure and laboratory parameters at admission to the hosting dialysis units were collected only in Poland ( $N = 161$ ). Since their last dialysis, 36.9% of patients had gained more than 3 kg of body weight, and 28.5% between 2 and 3 kg. Thirteen patients (8%) presented with a blood pressure  $\geq 180/100$  mmHg, an additional eight patients (5%) had systolic values above 180 mmHg, with diastolic values below 100 mmHg.

Detailed data on laboratory parameters are provided in Table 5. A substantial number of patients were admitted with marked anemia or hyperkalemia.

War-related injury (right shin hematoma) was reported in only one patient.

## DISCUSSION

The Russian invasion of Ukraine forced some patients treated by dialysis to migrate to other places in Ukraine or abroad. The RDRTF established by the ERA coordinated comprehensive support for kidney patients in Ukraine. The whole spectrum of its mission and activities to date has been described elsewhere [11]. In this study we present results of an international survey conducted by the ERA Task Force which analyzed the distribution, preparedness and management of adult Ukrainian dialysis patients who migrated outside Ukraine since the Russian invasion.

Our survey documented that around 6% of all patients dialyzed in Ukraine migrated outside the country. This number may be underestimated because of incomplete distribution and/or response rates of the questionnaire, migration to non-included countries and patients who died before being captured in the survey. On the other hand, data response rates in Poland and within the FMC network were high, suggesting that we captured a large proportion of the total number of displaced patients. However, complete data on migration within Ukraine, mostly to the western part, or of mortality

among Ukrainian dialysis patients is not available. Some data on relocation has been described elsewhere, including patients dialyzed in FMC centers in Chernihiv and Kharkiv who were moved to centers in Lviv and Cherkasy [12].

The proportion of dialysis patients who left the country was lower than one might have expected and a lot lower compared with migration of the general population [602 of 9648 (~6%) vs 7.5 million of 43.8 million (~17%)], likely reflecting enhanced vulnerability and limited mobility for various reasons of dialysis patients.

The mean age of dialysis patients fleeing Ukraine to foreign countries was  $48.1 \pm 13.4$  years, which is markedly lower than the mean age in Europe of 62.1 years [13]. Precise data on the age of dialysis patients in Ukraine before the war is not available, but is on the mean age of all KRT patients together (on dialysis and with a kidney transplant), which was 53.4 years, and therefore at baseline was lower than the corresponding age in other European countries [14]. As the mean age of those who migrated was lower than the mean before the war, it is possible that mostly younger and less vulnerable dialysis patients were able to migrate outside Ukraine. This finding is in line with a report on Syrian hemodialysis refugees in Turkey by Gursu *et al.* [15].

Consistent with this is the observation that the proportion of patients with kidney failure from diabetic nephropathy (often associated with multiple comorbidities) among the displaced population was lower than that in the total Ukrainian dialysis population (12.7% vs 15.7%) [14].

It may follow, that if younger and less vulnerable dialysis patients left the country, the general health condition of the remaining cohort was worse and more demanding at a time when resources were suboptimal. This should be taken into account in preparation of crisis management plans, and the need for different emergency preparedness strategies to minimize risk of life-threatening complications both in those remaining in the conflict zones and those fleeing war-affected regions [16–19].

As expected, the majority of patients migrated to the neighboring countries, i.e. to Poland and Slovakia. Only about one-fourth of patients decided to subsequently leave the reporting dialysis center and to move to another city or country. These results may suggest that about three-fourths of the patients had directly migrated to a previously predetermined center of their choice, or that they were satisfied with the location where they arrived.

An important clinical finding is that during the migration almost 30% of patients remained without dialysis for 4 days or longer. This highlights the necessity for close cooperation between dialysis centers in the conflict area and in neighboring countries, and development of transit dialysis options and facilitation of transportation to minimize delays in accessing therapy [20].

The main clinical challenges on presentation were hyperkalemia, fluid overload and severe anemia which could mainly be attributed to longer intervals between dialysis sessions. Severe anemia and fluid overload were among the main reasons for hospitalization upon arrival in the host center. Low hemoglobin concentrations were likely associated with fluid overload and missed ESA administration or lack of its availability before migration.

The proportion of displaced patients treated with ESA, active vitamin D and phosphate binders was rather low compared with the average European dialysis population [21, 22]. Possible reasons for this may be that the dialysis refugees from Ukraine were younger than average dialysis patients in Europe, and therefore had fewer comorbidities. Given that all patients from Ukraine received full healthcare coverage by the host healthcare system, guaranteeing standard therapy, including for kidney failure and listing for kidney transplantation if appropriate, following the international clinical and ethics guidance [23], it is unlikely that these therapies were withheld from the migrant patients.

A key challenge to optimal treatment was the lack of complete medical records in most patients, with an absence of all documentation in almost one in five patients. These findings support the importance of developing strategies of preparedness in the dialysis community. Another major problem was the language barrier, underlining the need to identify ahead of time translation resources for languages spoken by an anticipated refugee patients [20]. Contrary to our finding of the importance of a language barrier, cultural rather than linguistic differences were identified as a barrier for optimal care in the prior international survey of the nephrology community on KRT for refugees [24]. Cultural diversity in that study was however greater than ours, and migrants had had the opportunity to live in the host country for a long time, and therefore language barriers may have diminished over time.

To the best of our knowledge, this is the first report describing the clinical characteristics, distribution, preparedness and management of adult Ukrainian refugees on dialysis and one of the few systematic analyses of displaced dialysis patients during an active conflict.

The study does however have some limitations. Not all centers invited to participate responded. In addition, clinical and laboratory characteristics of these patients were requested

only from dialysis centers in Poland. Finally, 51.8% of patients were treated in the FMC centers and only aggregated data were obtainable due to internal privacy regulations. Despite these limitations, a major strength of this study is that it is one of few reports providing important information on migrant dialysis patients during conflicts and emergencies.

In summary, war always has a serious impact on the civilian population, especially those suffering from chronic diseases [25]. People with kidney disease are especially vulnerable given the complexity and technicality of kidney care [26]. Many European and other countries organized humanitarian aid for Ukraine. The ERA participated in this process after establishing the RDRTF. One of the many activities of this group [11] was conducting the present international survey. The results of this study will be helpful in understanding the scale of migration of dialysis patients outside Ukraine, their clinical characteristics, as well as medical and social needs, and may be relevant for patients in similar conflicts occurring elsewhere, and to support global recognition of this small but highly vulnerable group of patients. In addition, these data may be useful for Ukrainian nephrology community and authorities as they plan support for the patients who remained in Ukraine, who migrated within the country and for those migrants who intend to return to Ukraine [11, 12].

## CONCLUSIONS

- 1 After the Russian invasion of Ukraine, a relatively small proportion of Ukrainian dialysis patients appears to have fled their country and the majority of them chose a country neighboring Ukraine as the place for dialysis.
- 2 Preparedness for displacement varied and was incomplete in most patients.
- 3 During their transfer almost 30% of the patients remained without dialysis for 4 days or longer. Clinical status upon arrival necessitated hospitalization in one in three patients.
- 4 Full reimbursement of treatment enabled the receiving centers to support patients conforming to standard of care, including waitlisting for transplantation, if appropriate.
- 5 Almost one-third of patients not continuing treatment in the host country intended to return to Ukraine; however, return of these patients should be only encouraged when safety and treatment possibilities are guaranteed in their native country.
- 6 Results from our survey may inform evidence-based policies and interventions to prepare for and respond to the special needs of vulnerable kidney failure populations during armed conflicts and other emergencies.

## SUPPLEMENTARY DATA

Supplementary data are available at *ndt* online.

## ACKNOWLEDGEMENTS

Members of the Renal Disaster Relief Task Force of the European Renal Association: Serhan Tuğlular—Chair; Kai-



Uwe Eckardt; Daniel Gallego; Dimitri Ivanov; Anna Klis; Yelena Loboda; Valerie Luyckx; Edita Noruišiene; Ionut Nistor; Ewa Pawłowicz-Szlarska; Rukshana Shroff; Andrej Škoberne; Stefano Stuard; Mehmet Sukru Sever; Raymond Vanholder; and Andrzej Wiecek. The authors would like to express their deep gratitude to all dialysis centers employees for participating in the survey. European Kidney Health Alliance is the recipient of support by the European Union in the context of the Annual Work Program 2022 on prevention of non-communicable diseases of EU4Health, topic ID EU4H-2022-PJ02, project # 101101220 PREVENTCKD.

## AUTHORS' CONTRIBUTIONS

E.P.-S., R.V., M.S.S., S.T., V.L., K-U.E and A.W. were responsible for the study concept and design. E.P.-S., I.N., A.S., S.S., R.G., E.N. and A.W. were responsible for the acquisition of data. E.P.-S., R.V., M.S.S. and A.W. were responsible for the analysis and interpretation of data. E.P.-S. and A.W. were responsible for drafting of the manuscript. R.V., M.S.S., S.T., V.L., K-U.E., D.G., D.I., I.N., R.S., A.S., S.S., R.G., E.N. and M.S. provided critical revision of the manuscript for important intellectual content. All authors approved the final version of the manuscript.

## FUNDING

No funds, grants, or other financial support was received for conducting this study.

## DATA AVAILABILITY STATEMENT

The data underlying this article will be shared on reasonable request to the corresponding author.

## CONFLICT OF INTEREST STATEMENT

All authors declare that they have no competing interests. The results presented in this article have not been published previously in whole or part, except in abstract format.

## REFERENCES

1. United Nations Regional Information Center for Western Europe official website. Available from: <https://unric.org/en/the-un-and-the-war-in-ukraine-key-information/> (4 January 2023, date last accessed).
2. United Nations High Commissioner for Refugees official website. Available from: <https://data.unhcr.org/en/situations/ukraine> (24 September 2022, date last accessed).
3. UN Office for the Coordination of Humanitarian Affairs official website. Available from: <https://reliefweb.int/report/ukraine/ukraine-internal-displacement-report-general-population-survey-round-3-17-april-2022> (15 December 2022, date last accessed).
4. Garry S, Checchi F. Armed conflict and public health: into the 21st century. *J Public Health* 2020;**42**:e287–98. <https://doi.org/10.1093/pubmed/fdz095>.
5. Kutner NG, Muntner P, Huang Y *et al*. Effect of Hurricane Katrina on the mortality of dialysis patients. *Kidney Int* 2009;**76**:60–766. <https://doi.org/10.1038/ki.2009.268>.
6. Sever MS, Ereğ E, Vanholder R *et al*. Features of chronic hemodialysis practice after the Marmara earthquake. *J Am Soc Nephrol* 2004;**15**:1071–6. <https://doi.org/10.1097/01.ASN.0000119145.40232.67>.

7. Sever MS, Vanholder R, Luyckx V *et al*. Armed conflicts and kidney patients: a consensus statement from the Renal Disaster Relief Task Force of the ERA. *Nephrol Dial Transplant* 2022;**38**:56–65. <https://doi.org/10.1093/ndt/gfac247>.
8. Kolesnyk M, Driianska V, Liksunova L *et al*. State Institution 'Institute of Nephrology of the National Academy of Medical Sciences of Ukraine': 20 years afterward. *Ukr J Nephrol Dial* 2021;**4**:3–9.
9. Council Implementing Decision (EU) 2022/382 of 4 March 2022 establishing the existence of a mass influx of displaced persons from Ukraine within the meaning of Article 5 of Directive 2001/55/EC, and having the effect of introducing temporary protection. Available from: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex:32022D0382> (26 February 2023, date last accessed).
10. Stepanova N, Kolesnyk M, Mithanib Z *et al*. Lifesaving care for patients with kidney failure during the war in Ukraine. *Clin J Am Soc Nephrol* 2022;**17**:1079–81. <https://doi.org/10.2215/CJN.04720422>.
11. Tuglular S, Luyckx V, Vanholder R *et al*. Lessons learnt during the war in Ukraine: a report from the Renal Disaster Relief Task force of ERA. *Nephrol Dial Transplant* 2023;gfa053. <https://doi.org/10.1093/ndt/gfa053>.
12. Novakivskyy V, Shurduk R, Grin I *et al*. War in Ukraine and dialysis treatment: human suffering and organizational challenges. *Clin Kidney J* 2023;**16**:676–83. <https://doi.org/10.1093/ckj/sfad003>.
13. ERA Registry: ERA Registry Annual Report 2020. Amsterdam, The Netherlands: Amsterdam UMC, Location AMC, Department of Medical Informatics, 2022.
14. ERA Registry: ERA Registry Annual Report 2019. Amsterdam, The Netherlands: Amsterdam UMC, Location AMC, Department of Medical Informatics, 2022.
15. Gursu M, Arici M, Ates K *et al*. Hemodialysis experience of a large group of Syrian refugees in Turkey: all patients deserve effective treatment. *Kidney Blood Press Res* 2019;**44**:43–51. <https://doi.org/10.1159/000498832>.
16. Murakami N, Siktel HB, Lucido D *et al*. Disaster preparedness and awareness of patients on hemodialysis after Hurricane Sandy. *Clin J Am Soc Nephrol* 2015;**10**:1389–96. <https://doi.org/10.2215/CJN.10181014>.
17. Kopp JB, Ball LK, Cohen A *et al*. Kidney patient care in disasters: emergency planning for patients and dialysis facilities. *Clin J Am Soc Nephrol* 2007;**2**:825–38. <https://doi.org/10.2215/CJN.01220307>.
18. Dent L, Finne K, Lurie N. Progress in emergency preparedness for dialysis care 10 years after Hurricane Katrina. *Am J Kidney Dis* 2015;**66**:742–4. <https://doi.org/10.1053/j.ajkd.2015.08.019>.
19. Morsch G, Chacko S, Randolph R. Disaster medicine: disaster preparedness planning. *FP Essent* 2019;**48**:7:23–6.
20. Van Biesen W, Vanholder R, Hernandez T *et al*. Caring for migrants and refugees with end-stage kidney disease in Europe. *Am J Kidney Dis* 2018;**71**:701–9.
21. Fuller DS, Robinson BM, Locatelli F *et al*. Patterns of erythropoiesis-stimulating agent use in European hemodialysis patients: the Dialysis Outcomes and Practice Patterns study. *Nephron* 2018;**140**:24–30. <https://doi.org/10.1159/000490202>.
22. Fernández-Martín JL, Carrero JJ, Benedik M *et al*. COSMOS: the dialysis scenario of CKD-MBD in Europe. *Nephrol Dial Transplant* 2013;**28**:1922–35. <https://doi.org/10.1093/ndt/gfs418>.
23. Martin DE, Fadhil RAS, Więcek A. Ethical aspects of kidney donation and transplantation for migrants. *Semin Nephrol* 2022;**42**:151271. <https://doi.org/10.1016/j.semnephrol.2022.07.005>.
24. Van Biesen W, Vanholder R, Vanderhaegen B *et al*. Renal replacement therapy for refugees with end-stage kidney disease: an international survey of the nephrological community. *Kidney Int Suppl* 2016;**6**:35–41.
25. Vanholder R, Gallego D, Sever MS. Wars and kidney patients: a statement by the European Kidney Health Alliance related to the Russian-Ukrainian conflict. *J Nephrol* 2022;**35**:377–80. <https://doi.org/10.1007/s40620-022-01301-4>.
26. Vanholder R, De Wegheleire A, Ivanov D *et al*. Continuing kidney care in conflicts. *Nat Rev Nephrol* 2022;**18**:479–80. <https://doi.org/10.1038/s41581-022-00588-7>.

Received: 2.4.2023; Editorial decision: 12.4.2023