

The prevalence of posture pathology in school-aged children (results of a population-based study with the use of the computerized photo-geometric program "Posture")

I. O. Afanasieva*^{1,A-D,F}, M. V. Khaitovych^{1,A,C,E,F}, V. S. Potaskalova^{1,A-C,F},
I. V. Andrushchenko^{1,2,A-C,F}, V. V. Yevminov^{3,A,B,F}, I. A. Kudelia^{3,A,B,F}

¹Bogomolets National Medical University, Kyiv, Ukraine, ²SI "Institute of Nuclear Medicine and Diagnostic Radiology of the NAMS of Ukraine", Kyiv, ³Yevminov Vertebral Therapeutic Center, Kyiv, Ukraine

A – research concept and design; B – collection and/or assembly of data; C – data analysis and interpretation; D – writing the article; E – critical revision of the article; F – final approval of the article

Pathology of posture in children remains currently unresolved problem. According to foreign studies in the general population, the occurrence of idiopathic scoliosis in adolescents with an angle above 10° according to Cobb accounts for 0.93 % to 12.00 %. In Ukraine, the prevalence of posture pathology is not known.

The aim – to study the prevalence of posture pathology among school-aged children.

Material and methods. 1922 schoolchildren of 6–17 years, residents of Kyiv (Ukraine), were examined with the computerized photo-geometric program "Posture" between 2017 and early 2019.

Results. Deviation of the spinal column from the midline in the frontal, sagittal plane (functional and organic disorders of posture) was observed in 75.3 % of children. Among them, signs of scoliotic spinal deformity were found in 72.5 % of children, in girls – 1.5 times more often. The angle of maximum deflection of the scoliotic arch from the median line was $\geq 5^\circ$ in 13.7 % of children with scoliosis and in 9.9 % of all the examined children.

Conclusions. Sensitivity (95 %) and specificity (85 %) of the computerized photo-geometric program "Posture" were determined, which points to the necessity to be applied in health care maintaining practice. The pathology of posture in schoolchildren of the city of Kyiv was determined in 75.3 %, among them the signs of scoliotic spinal deformity were noted in 72.5 %. The scoliosis incidence with a deflection angle of spinal deformity $\geq 5^\circ$ was 9.9 %. It attests to the fact that it is necessary to develop medical and social programs for the prevention and treatment of posture pathology in children.

Key words:

screening, schoolchildren, scoliosis, incidence.

Zaporozhye medical journal
2020; 22 (3), 389-394

*E-mail:
afanasevaina@meta.ua

Поширеність патології постави в учнів (результати популяційного дослідження з використанням комп'ютеризованої фото-геометричної програми «Постава»)

I. A. Афанасьєва, М. В. Хайтович, В. С. Потаскалова, І. В. Андрущенко, В. В. Євмінов, І. А. Куделя

Патологія постави в дітей залишається невирішеною проблемою. За даними закордонних дослідників, у загальній популяції частота діагностики в підлітків ідіопатичного сколіозу з кутом понад 10° за Cobb становить від 0,93 % до 12,00 %. В Україні поширеність патології постави невідома.

Мета роботи – вивчити поширеність патології постави в дітей шкільного віку.

Матеріали та методи. За період із 2017 до початку 2019 року, використовуючи комп'ютеризовану фото-геометричну програму «Постава», обстежили 1922 школярі віком 6–17 років, які постійно проживають у м. Києві (Україна).

Результати. Відхилення хребта від серединної лінії у фронтальній, сагітальній площині (функціональні й органічні порушення постави) спостерігали у 75,3 % дітей. Ознаки сколіотичної деформації хребта виявили у 72,5 % дітей, у дівчат у 1,5 рази частіше. Кут максимального відхилення сколіотичної дуги від серединної лінії становив $\geq 5^\circ$ у 13,7 % дітей зі сколіозом і 9,9 % усіх обстежених.

Висновки. Визначили чутливість (95 %) і специфічність (85 %) комп'ютеризованої фото-геометричної програми «Постава», що вказує на необхідність її широкого впровадження у практику охорони здоров'я. Патологія постави у школярів Києва визначена у 75,3 %, ознаки сколіотичної деформації хребта виявили в 72,5 %. Частота діагностики сколіозу з кутом відхилення деформації хребта $\geq 5^\circ$ становила 9,9 %. Результати свідчать про необхідність розроблення медико-соціальних програм профілактики та лікування патології постави в дітей.

Ключові слова:

скринінг, школярі, сколіоз, поширеність.

Запорізький медичний журнал.
2020. Т. 22, № 3(120).
С. 389-394

Распространенность патологии осанки у школьников (результаты популяционного исследования с использованием компьютеризированной фото-геометрической программы «Осанка»)

И. А. Афанасьева, Н. В. Хайтович, В. С. Потаскалова, И. В. Андрущенко, В. В. Евминов, И. А. Куделя

Патология осанки у детей остается нерешенной проблемой. По данным зарубежных исследований, в общей популяции частота встречаемости у подростков идиопатического сколиоза с углом более 10° по Cobb составляет от 0,93 % до 12,00 %. В Украине распространенность патологии осанки неизвестна.

Ключевые слова:

скрининг, школьники, сколиоз, распространенность.

Запорожский медицинский журнал.
2020. Т. 22, № 3(120).
С. 389-394

Цель работы – изучить распространенность патологии осанки среди детей школьного возраста.

Материалы и методы. За период с 2017 до начала 2019 года с использованием компьютеризированной фото-геометрической программы «Осанка» обследовали 1922 школьников в возрасте 6–17 лет, постоянно проживающих в г. Киеве (Украина).

Результаты. Отклонение позвоночника от срединной линии во фронтальной, сагиттальной плоскости (функциональные и органические нарушения осанки) отмечено у 75,3 % детей. Признаки сколиотической деформации позвоночника диагностированы у 72,5 % детей, у девочек в 1,5 раза чаще. Угол максимального отклонения сколиотической дуги от срединной линии составлял $\geq 5^\circ$ у 13,7 % детей со сколиозом и 9,9 % всех обследованных детей.

Выводы. Определена чувствительность (95 %) и специфичность (85 %) компьютеризированной фото-геометрической программы «Осанка», что указывает на необходимость ее широкого внедрения в практику охраны здоровья. Патология осанки у школьников города Киева установлена у 75,3 %, признаки сколиотической деформации позвоночника отмечены у 72,5 %. Частота встречаемости сколиоза с углом отклонения деформации позвоночника $\geq 5^\circ$ составила 9,9 %. Результаты свидетельствуют о необходимости разработки медико-социальных программ профилактики и лечения патологии осанки у детей.

Pathology of posture in children includes persistent (organic) and functional curvatures of the spinal column and remains, until recently, an unresolved problem. Children usually visit to a doctor with scoliosis II or higher, when obvious signs of scoliosis are already visualized at the simple examination of a child. Television, video games, a sedentary lifestyle, smartphones and tablets as well as a lack of regular physical activity are the most common risk factors for posture curvatures in children [1,2].

Approximately 10 % of these diagnosed cases require conservative treatment, and approximately 0.1–0.3 % require urgent correction of the deformity [3].

Therefore, it is so important to diagnose posture pathology in children timely [2,4,5]. The data on the epidemiology of this pathology are very contradictory.

Aim

The aim of our work was to study the prevalence of posture pathology in children of school-age applying the screening method of the computerized photo-geometric program “Posture” in Kyiv (Ukraine).

Material and methods

The study included 1922 schoolchildren residents of Kyiv (Ukraine) from various schools, at the ages from 6 to 17 years, between 2017 and early 2019.

All children were examined with computerized photo-geometric program “Posture”. Three age groups were defined: Group I – 676 (35.2 %) children of 6–9 years; Group II – 546 (28.4 %) children of 10–12 years; Group III – 700 (36.4 %) adolescents of 13–17 years.

Examination rules:

- The undress child wearing the underwear and socks. Longer hair of the girls examined should be pinned up so as not to cover the earlobes or neck contour.
- The initial position of the child: upright position, legs closed, the child is relaxed.

- Placement of control points in the frontal plane (acromions (determines shoulder misalignment), shoulder blades’ angles, waist triangles, iliac bones’ ridges, gluteal folds, popliteal fossae) and in the sagittal plane (tragus, mid-shoulder, hip joint projection, mid-knee joint, ankle).

- The place of photographing should be provided with a plumb line.

- The child should be photographed in its original position in 2 projections.

- Computerized photo processing after “Posture” program (using control points on the child’s body set by a specialist, skew angles are calculated (shoulders, shoulder blades, pelvis, etc.), the presence, value in degrees, location of the spinal curvature) are determined.

- Questionnaire filling: age, height, weight, grade, type and style of school bag carrying.

- Parents are given a card with processed photos and examination results as well as recommendations.

Examination and interpretation of the results obtained by the photo-geometric method “Posture” took 5 minutes. The photos were included in assessment forms for postural analysis and identification of changes in the frontal plane, sagittal plane, such as kyphosis, lordosis, hypokyphosis, hypolordosis, and anterior head tilt.

The sensitivity was define as $Se = A / (A + C)$, where A is the proportion of true positive test results; A + C is the total number of true scoliosis patients. The specificity was define as $Sp = D / (B + D)$, where D is the proportion of true negative test results among patients without spinal deformity; B + D – diagnostic information value [6].

Results

The sex-specific characteristics of schoolchildren included in our study are presented in Fig. 1 and 2.

Among 1922 examined persons, it was found that the deviation of the spine in sagittal and frontal plane (along the spinous processes) was observed in 1447 (75.3 %) children; accordingly, the rest 475 had physiological posture, the incidence of which decreased with the child’s age (Table 1).

Table 1. The prevalence of posture pathology in children of school-age

| Posture | Age | | | | | | | |
|---------|--------------------|----|----------------------|----|----------------------|----|-----------------|------|
| | 6–9 years, n = 676 | | 10–12 years, n = 546 | | 13–17 years, n = 700 | | Total, n = 1922 | |
| | abs | % | abs | % | abs | % | abs | % |
| N | 277 | 41 | 93 | 17 | 105 | 15 | 475 | 24.7 |
| PD | 399 | 59 | 453 | 83 | 595 | 85 | 1447 | 75.3 |

N: physiological posture; PD: postural disorder.




Fig. 1. Sex ratio of 1922 children.




Fig. 2. Age distribution of 1922 children.

Table 2. The prevalence of scoliosis in children at different ages

| Scoliosis | Age, n = 1922 | | | | | |
|-------------------------------------|--------------------|-----|----------------------|------|----------------------|-----|
| | 6-9 years, n = 676 | | 10-12 years, n = 546 | | 13-17 years, n = 700 | |
| | abs | % | abs | % | abs | % |
| Cobb angle $\geq 20^\circ$, n = 80 | 13 | 1.9 | 16 | 2.9 | 51 | 7.3 |
| SC angle $\geq 5^\circ$, n = 191 | 27 | 4.0 | 59 | 10.8 | 105 | 15 |

SC: spinal curvature in the frontal plane.

According to the study results, scoliotic disease was 1.5 times more often in girls.

Signs of scoliotic spinal deformity were detected in 1393 (72.5 %) children, kyphotic curvature – in 54, combined disorders were detected in 88 schoolchildren.

During data analysis, which were obtained with computerized photo-geometric "Posture" method, the angle of scoliotic arc maximum deflection from the median line was more than $\geq 5^\circ$ in 191 (13.7 %) children with scoliosis and in 190 (9.9 %) of all the examined children (Fig. 3).

Children with scoliosis were proposed X-ray investigation in an upright position. According to X-ray images, scoliosis of varying severity was detected in 80 children, in 5 of them – S-shaped scoliosis of III-IV stage of thoracolumbar area. The study results demonstrated that the prevalence of scoliosis Cobb angle of $\geq 20^\circ$ increased with the children age (Table 2).

Discussion

Deviation of the spinal column from the median line along the spinous processes in the frontal and sagittal planes (functional and organic disorders of posture) was observed in 75.3 % of children. Among them, signs of scoliotic spinal deformity were detected in 72.5 % of children, in girls 1.5 times more often. In 13.7 % of children with scoliosis and in 9.9 % of all the examined children, the angle of maximum deviation of the scoliotic arc from the midline was $\geq 5^\circ$.

According to epidemiological investigations in other countries, in the general population, the frequency of idiopathic scoliosis occurrence in adolescents with an angle above 10° according to Cobb ranges from 0.93 % to 12.00 %. In about 20 % of cases, scoliosis is secondary to another pathological process. The remaining 80 % are cases of idiopathic scoliosis [7-11].




Fig. 3. Computerized photo-geometric program "Posture": measurement of the spinal deviation angle in a child.

Among 79 122 schoolchildren in the city of Wuxi, eastern China, adolescent idiopathic scoliosis was found in 2.4 %, and in the Shanghai area, among 5327 investigated secondary school students (grades 6–8), scoliosis was diagnosed by screening method in 520 children (9.76 %) and spinal X-ray was performed in 102 of them (33.9 %) [12,13].

Ueno M., et al. performed school screenings for more than 250 000 children aged 11–14 years from 2003 to 2007 in the city of Tokyo. Scoliosis was confirmed in 3424 children by radiographic examination, and false-positive results were found (Cobb 0) in 39 cases [14].

2822 schoolchildren were examined in Spain (mean age 8.5 ± 1.8 years), where the number of patients with scoliosis was 1023 (36.3 %). Rib hump was twice common in girls. The risk of hump developing increased with age by 1.12 times. Children wearing school backpacks had a lower risk of scoliosis progression [15].

966 schoolchildren aged 10–16 years were examined in Southern Portugal. The results of the study demonstrated that scoliosis had been present in 41 (4.2 %) children. The highest prevalence was observed among girls (4.5 %), especially at delayed menarche (8.6 %) [16].

In some Polish cities, there are preventive programs aimed at detection of spinal dysfunctions and disorders in children and adolescents. Our study participants were children aged between 6 and 18 years with permanent residence in Poznan. The study enrolled 1.325 boys and 1.355 girls. In the whole study population, the incidence of scoliotic posture, assessed according to the chart of postural dysfunctions by Dega, amounted to 14.6 % in boys and 17.0 % in girls. There were no significant differences between the boys and the girls in particular age groups. It was found that scoliotic posture both in boys and girls ($P < 0.05$) was associated with age (the correlation was significantly different from zero and positive), and the proportions of students with scoliotic posture rose with age. Scoliometer examination revealed that the incidence of scoliotic posture was more frequent among girls (25.7 %) than among boys (22.2 %). The differences were statistically significant. The boys subjected to the study showed a correlation ($r = 0.12$) significantly different from zero and positive, in the group of tested girls, the correlation ($r = 0.14$) was significantly different from zero and positive [17].

The screening program in Hong Kong included 2242 children aged 8–16 years. Sensitivity and specificity (93.8 % and 99.2 %) and predictive value (81.0 %) were the highest and false-positive indexes were the lowest (0.8 %; 6.2 % false-negative) [18].

Chaves P. J. et al. examined 117 schoolchildren in Brazil and stated that high incidence of postural changes was observed in the studied population. About 56 % ($n = 14$) presented some type of head alteration and the other 44 % ($n = 11$) presented no head position changes. Among the postural alteration in the alignment of the head, 12 % represented head protrusion and 44 % represented head tilt to the right or left. 64 % represented a shoulder elevation and 24 % had a shoulder protrusion. Other postural alterations also were observed in the studied population, and 67.27 % ($n = 74$) had pelvic alterations, such as anteversion and pelvic retroversion [19].

The scoliosis incidence among adolescents in Malaysia was 2.55 % [20].

In the United States of America, according to J. Dunn, 2017, the cumulative scoliosis incidence was 1.8 % (reliability 95 %): the interval [CI] of 1.2–2.3 for curvature of more than 10° ; 1.0 % (95 % CI, 0.6–1.5) for deformation of at least 20° and 0.4 % (95 % CI; 0.1–0.6) for curvature of 40° [17,21].

Referring to the literature data [6], we have established sensitivity and specificity of the computerized photo-geometric program "Posture", which has indicators 95 % and 85 %, respectively. Diagnostic information content of this method is 0.9.

The spine deviation angle, equal to 5° and determined by a scoliometer, is a criterion for determining the Cobb angle equal to 20° in computer analyzes [22–24]. In our studies, scoliosis of the II degree and higher was observed in 9.9 % of 1922 schoolchildren in Kyiv, respectively, 13.7 % of children had all types of scoliosis, which exceeded the data of modern researchers of the world [2,7,20].

When comparing the data of modern foreign literature [12,13,20,23], the incidence indicators of scoliosis (72.5 %) and postural disorder (75.0 %) in school-aged children coincided.

Such a high prevalence of posture pathology in children is probably because of the lack of medical examination, a change in the lifestyle of modern children (long "monitor" time, using of gadgets, low physical activity, etc.) [25].

Quite possible that the personalized approach to the treatment may contribute to regression and/or compensation, reducing the risk of scoliotic disease further progression [26–30]. Treatment result was provided with an integrated approach using exercises, physiotherapy, corset wearing and, possibly, patient compliance [30–32].

According to the Australian Association of Physiotherapists, the correct balance between screen time and physical activity should be achieved for the prevention and treatment of posture pathology [33]. Therefore, it is so important to diagnose pathology at the early stages.

Conclusions

Sensitivity (95 %) and specificity (85 %) of the computerized photo-geometric program "Posture" were determined, which point to the necessity to be applied in health care maintaining practice.

Posture disorders in schoolchildren of Kyiv were diagnosed in more than 75 % among them signs of scoliotic spinal deformity were found in 72.5 %. The scoliosis incidence with a deflection angle of spinal deformity $\geq 5^\circ$ was 9.9 %.

It attests to the fact that it is necessary to develop medical and social programs for the prevention and treatment of posture pathology in children.

Practical recommendations

1. It is recommended to provide photo-geometric screening to school-aged children in order to detect posture disorders.
2. Detection of posture disorders in children will prevent the development of scoliosis of the II degree and higher.

Conflicts of interest: authors have no conflict of interest to declare.
Конфлікт інтересів: відсутній.

Надійшла до редакції / Received: 28.08.2019

Після доопрацювання / Revised: 11.11.2019

Прийнято до друку / Accepted: 29.11.2019

Information about authors:

Afanasiyeva I. O., MD, PhD, Associate Professor, Department of Clinical Pharmacology and Clinical Pharmacy, Bogomolets National Medical University, Kyiv, Ukraine.
 Khaitovych M. V., MD, PhD, DSc, Professor, Head of the Department of Clinical Pharmacology and Clinical Pharmacy, Bogomolets National Medical University, Kyiv, Ukraine.
 Potaskalova V. S., MD, PhD, Associate Professor, Department of Clinical Pharmacology and Clinical Pharmacy, Bogomolets National Medical University, Kyiv, Ukraine.
 Andrushchenko I. V., MD, PhD, Senior Researcher, SI "Institute of Nuclear Medicine and Diagnostic Radiology of the NAMS of Ukraine", Scientific Secretary, Kyiv.
 ORCID ID: [0000-0001-8199-9941](https://orcid.org/0000-0001-8199-9941)
 Yevminov V. V., Head of Evminov Vertebral Therapeutic Center, Kyiv, Ukraine.
 Kudelia I. O., Master in Physical Rehabilitation, National University of Ukraine on Physical Education and Sport, Kyiv, Ukraine.

Відомості про авторів:

Афанасьєва І. О., канд. мед. наук, доцент каф. клінічної фармакології та клінічної фармації, Національний медичний університет імені О. О. Богомольця, м. Київ, Україна.
 Хайтович М. В., д-р мед. наук, зав. каф. клінічної фармакології та клінічної фармації, Національний медичний університет імені О. О. Богомольця, м. Київ, Україна.
 Потаскалова В. С., канд. мед. наук, доцент каф. клінічної фармакології та клінічної фармації, Національний медичний університет імені О. О. Богомольця, м. Київ, Україна.
 Андрущенко І. В., канд. мед. наук, старший науковий співробітник, ДУ «Інститут ядерної медицини та променевої діагностики НАМН України», вчений секретар, м. Київ.
 Євмінов В. В., керівник Вертебрально-оздоровчого центру Євмінова, м. Київ, Україна.
 Куделя І. О., магістр із фізичної реабілітації, Національний університет фізичного виховання і спорту України, м. Київ.

Сведения об авторах:

Афанасьева И. А., канд. мед. наук, доцент каф. клинической фармакологии и клинической фармации, Национальный медицинский университет имени А. А. Богомольца, г. Киев, Украина.
 Хайтович Н. В., д-р мед. наук, зав. каф. клинической фармакологии и клинической фармации, Национальный медицинский университет имени А. А. Богомольца, г. Киев, Украина.
 Потаскалова В. С., канд. мед. наук, доцент каф. клинической фармакологии и клинической фармации, Национальный медицинский университет имени А. А. Богомольца, г. Киев, Украина.
 Андрущенко И. В., канд. мед. наук, старший научный сотрудник, ГУ «Институт ядерной медицины и лучевой диагностики НАМН Украины», ученый секретарь, г. Киев.
 Евминов В. В., руководитель Вертебрально-оздоровительного центра Евминова, г. Киев, Украина.
 Куделя И. А., магистр по физической реабилитации, Национальный университет физического воспитания и спорта Украины, г. Киев.

References

- Karimi, M. T., & Rabczuk, T. J. (2018). Scoliosis conservative treatment: A review of literature. *Journal of Craniovertebral Junction and Spine*, 9(1), 3-8. https://doi.org/10.4103/jcvjs.JCVJS_39_17
- Negrini, S., Donzelli, S., Aulisa, A. G., Czuprowski, D., Schreiber, S., de Mauroy, J. C., Diers, H., Grivas, T. B., Knott, P., Kotwicki, T., Lebel, A., Marti, C., Maruyama, T., O'Brien, J., Price, N., Parent, E., Rigo, M., Romano, M., Stikeleather, L., ... Zaina, F. (2018). 2016 SOSORT guidelines: orthopaedic and rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis and Spinal Disorders*, 13, Article 3. <https://doi.org/10.1186/s13013-017-0145-8>
- Bozzola, E., Spina, G., Ruggiero, M., Memo, L., Agostiniani, R., Bozzola, M., Corsello, G., & Villani, A. (2018). Media devices in pre-school children: the recommendations of the Italian pediatric society. *Italian Journal of Pediatrics*, 44(1), Article 69. <https://doi.org/10.1186/s13052-018-0508-7>
- Jada, A., Mackel, C. E., Hwang, S. W., Samdani, A. F., Stephen, J. H., Bennett, J. T., & Baaj, A. A. (2017). Evaluation and management of adolescent idiopathic scoliosis: a review. *Neurosurgical Focus*, 43(4), Article E2. <https://doi.org/10.3171/2017.7.focus17297>
- Cobetto, N., Aubin, C. -E., & Parent, S. (2018). Surgical Planning and Follow-up of Anterior Vertebral Body Growth Modulation in Pediatric Idiopathic Scoliosis Using a Patient-Specific Finite Element Model Integrating Growth Modulation. *Spine Deformity*, 6(4), 344-350. <https://doi.org/10.1016/j.jspd.2017.11.006>
- Fletcher, R. H., Fletcher, S. W., & Wagner, E. H. (1998). *Klinicheskaya epidemiologiya. Osnovy dokazatel'noi meditsiny [Clinical epidemiology. The essentials]* (3rd ed.). Mediya Sfera. [in Russian].
- Ciaccia, M. C. C., Castro, J. S., Rahal, M. A., Penatti, B. S., Selegatto, I. B., Giampietro, J. L. M., & Rullo, V. E. V. (2017). Prevalence of scoliosis in public elementary school students. *Revista Paulista de Pediatria*, 35(2), 191-198. <https://doi.org/10.1590/1984-0462/2017/35/2/00008>
- Mahabadi, E. A., Ghandehari, H., Mahdavi, S. M., Vahid Tari, S. H., Sotoudeh, A. S., & Safdari, F. (2015). Adolescent Idiopathic Scoliosis: Males Versus Females. *Shafa Orthopedic Journal*, 2(2), Article e901. <https://doi.org/10.17795/scj-901>
- Stępień, A., Guzek, K., Czubak, J., Surowińska, J., Stępowka, J., & Kiebzak, W. (2017). Early childhood anomalies of the hip occur at a similar frequency in patients with idiopathic scoliosis and in healthy individuals – questionnaire for parents. *Advances in Rehabilitation*, 4(4), 5-18. <https://doi.org/10.1515/rehab-2015-0075>
- Bozkurt, S., Kayalar, G., Tezel, N., Güler, T., Kesikburun, B., Denizli, M., Tan, S., & Yilmaz, H. (2019). Hypermobility Frequency in School Children: Relationship With Idiopathic Scoliosis, Age, Sex and Musculoskeletal Problems. *Archives of rheumatology*, 34(3), 268-273. <https://doi.org/10.5606/ArchRheumatol.2019.7181>
- Grabala, P. (2019). Idiopathic scoliosis in children and adolescents – a literature review. *Journal of Education, Health and Sport*, 9(6), 212-217. <https://doi.org/10.5281/zenodo.3241930>
- Zheng, Y., Dang, Y., Wu, X., Yang, Y., Reinhardt, J., He, C., & Wong, M. (2017). Epidemiological study of adolescent idiopathic scoliosis in Eastern China. *Journal of Rehabilitation Medicine*, 49(6), 512-519. <https://doi.org/10.2340/16501977-2240>
- Guo, Y., Jiang, Q., Tanimoto, T., Kami, M., Luo, C., Leppold, C., Nishimura, K., He, Y., Kato, S., & Ding, X. (2017). Low hospital referral rates of school scoliosis screening positives in an urban district of mainland China. *Medicine*, 96(14), Article e6481. <https://doi.org/10.1097/md.00000000000006481>
- Cheng, J. C., Castelein, R. M., Chu, W. C., Danielsson, A. J., Dobbs, M. B., Grivas, T. B., Gurnett, C. A., Luk, K. D., Moreau, A., Newton, P. O., Stokes, I. A., Weinstein, S. L., & Burwell, R. G. (2015). Adolescent idiopathic scoliosis. *Nature Reviews Disease Primers*, 1, Article 15030. <https://doi.org/10.1038/nrdp.2015.30>
- Ortega, Z. F., Fernández, S. M., Fernández, G. R., Schyke, C. E. J., & Morales, Z. L. (2014). Predictors of scoliosis in school-aged children. *Gaceta medica de Mexico*, 150(6), 533-539. [in Spanish].
- Minghelli, B., Nunes, C., & Oliveira, R. (2014). Prevalence of scoliosis in southern Portugal adolescents. *Pediatric endocrinology reviews*, 11(4), 374-382.
- Maciałczyk-Paprocka, K., Stawińska-Witoszyńska, B., Kotwicki, T., Sowińska, A., Krzyżaniak, A., Walkowiak, J., & Krzywińska-Wiewiorowska, M. (2017). Prevalence of incorrect body posture in children and adolescents with overweight and obesity. *European Journal of Pediatrics*, 176(5), 563-572. <https://doi.org/10.1007/s00431-017-2873-4>
- Dunn, J., Henrikson, N. B., Morrison, C. C., Nguyen, M., Blasi, P. R., & Lin, J. S. (Eds.). (2018). *Screening for Adolescent Idiopathic Scoliosis: A Systematic Evidence Review for the U.S. Preventive Services Task Force* [Internet]. Agency for Healthcare Research and Quality. Evidence Synthesis, (156). <https://www.ncbi.nlm.nih.gov/books/NBK493364/>
- Rusnák, R., Kolarová, M., Aštaryová, I., & Kutíš, P. (2019). Screening and Early Identification of Spinal Deformities and Posture in 311 Children: Results from 16 Districts in Slovakia. *Rehabilitation Research and Practice*, 2019, Article 4758386. <https://doi.org/10.1155/2019/4758386>
- Deepak, A. S., Ong, J. Y., Choon, D., Lee, C. K., Chiu, C. K., Chan, C., & Kwan, M. K. (2017). The Clinical Effectiveness of School Screening Programme for Idiopathic Scoliosis in Malaysia. *Malaysian Orthopaedic Journal*, 11(1), 41-46. <https://doi.org/10.5704/moj.1703.018>
- Dunn, J., Henrikson, N. B., Morrison, C. C., Blasi, P. R., Nguyen, M., & Lin, J. S. (2018). Screening for Adolescent Idiopathic Scoliosis. *Journal of the American Medical Association*, 319(2), 173-187. <https://doi.org/10.1001/jama.2017.11669>

- [22] Afanasieva, I. O., & Yevminov, V. V. (2019). Sposib diahnostryky oporno-rukhevoho aparatu u ditei [Method for diagnosing musculoskeletal system in children] (Ukraine. Patent No. 135078). Ukraina. Derzhavna sluzhba intelektualnoi vlasnosti Ukrainy. <https://base.uipv.org/search/NV/search.php?action=viewdetails&IdClaim=259359> [in Ukrainian].
- [23] do Rosário, J. L. P. (2014). Photographic analysis of human posture: A literature review. *Journal of Bodywork and Movement Therapies*, 18(1), 56-61. <https://doi.org/10.1016/j.jbmt.2013.05.008>
- [24] Kokabu, T., Kawakami, N., Uno, K., Kotani, T., Suzuki, T., Abe, Y., Maeda, K., Inage, F., Ito, Y. M., Iwasaki, N., & Sudo, H. (2019). Three-dimensional depth sensor imaging to identify adolescent idiopathic scoliosis: a prospective multicenter cohort study. *Scientific Reports*, 9(1), Article 9678. <https://doi.org/10.1038/s41598-019-46246-0>
- [25] Plaszewski, M., Kotwicki, T., Chwata, W., Terech, J., & Cieśliński, I. (2015). Study protocol and overview of the literature on long-term health and quality of life outcomes in patients treated in adolescence for scoliosis with therapeutic exercises. *Journal of Back and Musculoskeletal Rehabilitation*, 28(3), 453-462. <https://doi.org/10.3233/bmr-140540>
- [26] Donzelli, S., Zaina, F., & Negrini, S. (2012). In defense of adolescents: They really do use braces for the hours prescribed, if good help is provided. Results from a prospective everyday clinic cohort using thermobrace. *Scoliosis*, 7(1), Article 12. <https://doi.org/10.1186/1748-7161-7-12>
- [27] Romano, M., Negrini, A., Parzini, S., Tavernaro, M., Zaina, F., Donzelli, S., & Negrini, S. (2015). SEAS (Scientific Exercises Approach to Scoliosis): a modern and effective evidence based approach to physiotherapeutic specific scoliosis exercises. *Scoliosis*, 10, Article 3. <https://doi.org/10.1186/s13013-014-0027-2>
- [28] Ko, J. Y., Suh, J. H., Kim, H., & Ryu, J. S. (2018). Proposal of a new exercise protocol for idiopathic scoliosis. *Medicine*, 97(49), Article e13336. <https://doi.org/10.1097/md.00000000000013336>
- [29] Burton, D. C., Carlson, B. B., Place, H. M., Fuller, J. E., Blanke, K., Cho, R., Fu, K. -M., Ganju, A., Heary, R., Herrera-Soto, J. A., Larson, A. N., Lavelle, W. F., Nelson, I. W., Vernengo-Lezica, A., & Verska, J. M. (2016). Results of the Scoliosis Research Society Morbidity and Mortality Database 2009–2012: A Report From the Morbidity and Mortality Committee. *Spine Deformity*, 4(5), 338-343. <https://doi.org/10.1016/j.jspd.2016.05.003>
- [30] Schreiber, S., Parent, E. C., Moez, E. K., Hedden, D. M., Hill, D., Moreau, M. J., Lou, E., Watkins, E. M., & Southon, S. C. (2015). The effect of Schroth exercises added to the standard of care on the quality of life and muscle endurance in adolescents with idiopathic scoliosis-an assessor and statistician blinded randomized controlled trial: «SOSORT 2015 Award Winner». *Scoliosis*, 10, Article 24. <https://doi.org/10.1186/s13013-015-0048-5>
- [31] Kuru, T., Yeldan, İ., Dereli, E. E., Özdiñçler, A. R., Dikici, F., & Çolak, İ. (2015). The efficacy of three-dimensional Schroth exercises in adolescent idiopathic scoliosis: a randomised controlled clinical trial. *Clinical Rehabilitation*, 30(2), 181-190. <https://doi.org/10.1177/0269215515575745>
- [32] Kulikov, A. G., Zaytseva, T. N., Pyzhevskaya, O. P., & Ivanova, E. R. (2016). Skolioz u detei: novye podkhody k resheniyu vazhnoi mediko-sotsial'noi problemy [Scoliosis in children: new approaches to the solution of important medical and social problems]. *Mediko-sotsial'naya ekspertiza i reabilitatsiya* 19(4), 178-181. [in Russian].
- [33] Chiang, H. (Ariel), & Liu, C. -H. (2016). Exploration of the associations of touch-screen tablet computer usage and musculoskeletal discomfort. *Work*, 53(4), 917-925. <https://doi.org/10.3233/wor-162274>