

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/340279620>

A review of the medicinal Lycopodiophyta of Ukraine

Article in Scripta Scientifica Pharmaceutica · December 2019

DOI: 10.14748/ssp.v6i1.6206

CITATIONS

0

READS

113

3 authors:



Tetyana Sergiivna Dvirna
M.G.Kholodny Botany Institute

69 PUBLICATIONS 87 CITATIONS

[SEE PROFILE](#)



Valentyna Minarchenko
M.G.Kholodny Botany Institute

36 PUBLICATIONS 26 CITATIONS

[SEE PROFILE](#)



Iryna Andriivna Tymchenko
M.G.Kholodny Botany Institute

51 PUBLICATIONS 39 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Morphologo-anatomical and resource characteristics of related species of medicinal plants of Ukraine [View project](#)



THREATENED MEDICINAL PLANTS OF UKRAINE: AN ASSESSMENT OF THE CURRENT PROTECTION STATUS [View project](#)

A REVIEW OF THE MEDICINAL LYCOPODIOPHYTA OF UKRAINE

Tetyana Dvirna¹, Valentyna Minarchenko², Iryna Tymchenko¹

¹*Department of Systematics and Floristics of Vascular Plants,*

M.G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine

²*Department of Pharmacognosy and Botany, M.G. Kholodny Institute of Botany,
National Academy of Sciences of Ukraine*

ABSTRACT

The publication presents the results of complex research and analysis of medicinal plain-like of flora of Ukraine. *Lycopodiophyta* of Ukraine includes four families: *Lycopodiaceae* (8 species – *Diphasiastrum alpinum* (L.) Holub, *D. complanatum* (L.) Holub, *D. isselieri* (Rouy) Holub, *D. tristachyum* (Pursh) Holub, *D. zeilleri* (Rouy) Holub, *Lycopodiella inundata* (L.) Holub, *Lycopodium annotinum* L., *L. clavatum* L.), *Huperziaceae* (1 – *Huperzia selago* (L.) Bernh. ex Schrank & C. Mart.), *Selaginellaceae* (2 – *Selaginella helvetica* (L.) Spring., *Selaginella selaginoides* (L.) C. Mart.) and *Isoëtaceae* (1 – *Isoëtes lacustris* L.). They have different contents of biologically active substances and all of these species are medicinal or semi-medicinal. The paper presents the characteristics of their distribution in Ukraine, resource importance, population structure, environmental protection status in Ukraine and Europe, main active substances, and medicinal properties.

Keywords: medicinal *Lycopodiophyta*, resources significance, protection, Ukraine

INTRODUCTION

Nowadays, medicinal plants attract attention because they are among the main sources of medical and preventive means of non-traditional and scientific modern medicine (1-5).

One of the oldest and most interesting groups of plants is the representatives of *Lycopodiophyta*, which achieved their greatest development in the late Paleozoic. Currently it is represented by a relatively small number of genera and species, which account

for only 1% of the modern species diversity of vascular plants (6-8).

Extinct lycopods had an important role in the formation of coal (6, 9) while modern species have phytogenic, industrial, decorative and medicinal value (1-6, 10-12).

Lycopods are able to form continuous carpets in the woods and participate in the formation of plant communities, which constitutes their phytogenic value. Spores of Lycopods are rich in fatty oils and aluminum, which is a reason for their use in pyrotechnics (for the manufacture of fireworks), metallurgy (for smoothness of surfaces at packed casting), in the manufacture of photos, where they are a source of green, blue and yellow colors. Plants are harvested for making bouquets and for various rituals, some species of *Selaginella* are indoor ornamental plants (6,9,10,13).

The medicinal properties of the plants were noted by the ancient Romans, who considered them the best remedy for eliminating dullness of the skin (2). Nowadays, spores (lycopodium) are used as a

Address for correspondence:

Tetyana Dvirna
M.G. Kholodny Institute of Botany
National Academy of Sciences of Ukraine
2 Tereshchenkivska St
01601 Kiev (Kyiv)
Ukraine
e-mail: dvirna_t@ukr.net

Received: April 8, 2019

Accepted: June 24 2019

powder (as a baby powder, for treatment of bedsores), and for pill coating, grass lycopsids are used as a diuretic, anti-rheumatic, anti-inflammatory, and anesthetic agent, in the treatment of mental illness and Alzheimer's disease; species of *Huperzia* are poisonous but used for the treatment of alcoholism and nicotine (1-3,5,11,14).

Lycopodiophyta of Ukraine includes four families: *Lycopodiaceae* (8 species – *Diphasiastrum alpinum* (L.) Holub, *D. complanatum* (L.) Holub, *D. is seleri* (Rouy) Holub, *D. tristachyum* (Pursh) Holub, *D. zeilleri* (Rouy) Holub, *Lycopodiella inundata* (L.) Holub, *Lycopodium annotinum* L., *L. clavatum* L.), *Huperziaceae* (1 – *Huperzia selago* (L.) Bernh. ex Schrank & C. Mart.), *Selaginellaceae* (2 – *Selaginella helvetica* (L.) Spring., *Selaginella selaginoides* (L.) C. Mart.) and *Isoetaceae* (1 – *Isoëtes lacustris* L.) (7,8) are medicinal or semi-medicinal.

Below we present a more detailed description of these species (Table 1).

AIM AND METHODS

The object of our study are the medicinal and semi-medicinal species of the *Lycopodiophyta* representatives of the flora of Ukraine. The basis of the work is the data obtained during the field research as a result of working on literary materials and maps, and herbarium collections of Ukraine gathered in the period 2010–2019.

RESULTS AND DISCUSSION

Lycopodiophyta species are of no economic value in Ukraine due to several reasons. The results of the study are presented in Table 1 in which the checklist of representatives of the studied department is given in alphabetical order. The name of each species is in Latin and English (9 species), presented are also family, distribution in Ukraine, resource value, population structure, conservation status in Ukraine and Europe, main active substances, and medicinal properties.

We analyzed the distribution of plant species in Ukraine and established that the majority – 42% (5 species) — is known from several localities, mainly growing in the Carpathians and Polissya (10,12,13,15,16).

The studied species of *Lycopodiophyta* have no resource values since most species (83% or 10 species) in the wild are not available.

The population structure of the species is unstable. However, 42% (5 species) are represented by several individuals, which is caused by a number of threats, but the main one is the anthropogenic factor (cutting, grazing, recreation, etc.) to which species are very sensitive (10,13).

All species are listed in the Red Book of Ukraine and require conservation measures. Most of them – 42% (5 species) — are vulnerable and only one species (*Lycopodium clavatum*) is regionally rare. *L. clavatum* and *Huperzia selago* are included in the European Red List, to the IUCN database with LC status, and to the European Red List of Medicinal Plants (13,17-19).

The pharmacological properties of the species emerge due to the chemical composition and the ratio of active ingredients. For medicinal purposes, spores and aerial parts of plants are used as raw material. The analysis of literary data enabled us to establish that model species that are taken as the basis for the research of pharmacological properties usually are *L. clavatum* and *H. selago*. It is caused by their chemical composition, which slightly differs from “other composition and ratio of components”, while the properties, uses, and dosage forms are similar (1,20-24).

The main common active substances of *Lycopodiophyta* are alkaloids (lycopodine, clavatin, lycopodine, etc.), flavonoids, triterpenoids, flavones, guperine, carotenoids, fatty oils (spores) (1-3,5,6,11,20-38).

The spores of Lycopods contain about 50% non-drying fatty oil and glycerides, stearic, oleic, arachidonic, palmitic and other acids, therefore, they are used as a powder for infants, for the treatment of eczema, adulthood, bedsores, wounds, burns (1,2,4-6,11).

Spores and herbs are used in liver disease, respiratory tract diseases, and as diuretic, antirheumatic, anticonvulsant and anti-inflammatory, antibacterial and antifungal, antiviral and antiemetic agents, etc., as well as for the treatment of psychological disorders, Alzheimer's disease, alcoholism and nicotine (1-5,11,38).

Table 1. The state and medicinal values of Lycopodiophyta of Ukraine

Latin Name	English Name	Family	Distribu-tion in Ukraine	Resource Value	Number of Individuals in Popula-tion	Protection Status in Ukraine	Protection Status (by Global and European Regional Assess-ment)	Medicinal Value
<i>Diphasiastrum alpinum</i> (L.) Holub	Alpine Clubmoss	Lycopodiaceae	R	0	F	RB, rare	-	Alkaloids – O-acetylpiclavolamine, O-acetylfawcettine, O-acetylfololine, lycodoline, clavolonne, flavonoids, triterpenoids. It has an anesthetic effect, used in hydrophobia; some alkaloids of the species have proven to be effective inhibitors of acetylcholinesterase and are potentially able to alleviate the symptoms of Alzheimer's disease. Spores used as a dusting powder.
<i>Diphasiastrum complanatum</i> (L.) Holub	Ground-cedar	Lycopodiaceae	R	0	F	RB, rare	-	Serratane triterpenoids, including serratane-3 α ,14 α ,15 α ,20 β ,21 β -heptol, 3 α ,20 β ,21 β -trihydroxyserrat-14-en-24-oic acid, 3 β ,20 β ,21 β -trihydroxyserrat-14-en-24-oic acid, 3 α ,20 β ,21 β -trihydroxy-16-oxoserrat-14-en-24-oic acid, and 16-oxolycianitin-29-yl E-4'-hydroxy-3'-methoxycinnamate, sterols, lycopadins D, E and complanadine E, β carotene, lipids, fatty oil; triterpenoids, lysine-derived alkaloids. It has an anesthetic, diuretic and sedative effect; used in the treatment of tachycardia, flu, migraine, spasmosis, liver disease, diarrhea, infertility, metabolic disorders, externally used in alopecia, baldness, dermatomycosis. Spores used as a dusting powder.
<i>Diphasiastrum isselieri</i> (Rouy) Holub	Issler's Clubmoss	Lycopodiaceae	L	0	I	RB, vulnerable	-	Lycopodine ($C_{16}H_{22}NO$); lysine-derived alkaloids. Ways to use in medicine aren't explored. Spores used for powders.

<i>Diphasiastrum tristachyum (Pursh) Holub</i>	Blue ground- cedar	Lycopodiaceae	L	0	I	RB, endangered	-	Lysine-derived alkaloids; lycopodine; alkaloid L 13-15, carotenoids, a small amount of nicotine, rare flavones: chrysoeriol, segin and tricin; triterpenoids; Lycodine type alkaloids, tetracyclic alkaloids, clavine alkaloid, huperzine A. Antifungal, antibacterial, antiviral properties; relieve spasms, increase urine flow, fever, inflammation; insecticidal. Disputes are used as a baby powder and in the treatment of bedsores; in pharmacy - for powder tablets.
<i>Diphasiastrum zeilleri (Rouy) Holub</i>	Zeller's clubmoss	Lycopodiaceae	L	0	I	RB, endangered	-	Lycodine type alkaloids, lycopodine alkaloids, tetracyclic alkaloids, clavine alkaloid, huperzine A. Antifungal, antibacterial, antiviral; relieve spasms, increase urine flow, estrogenic; reduce pain, fever, inflammation; insecticidal.
<i>Lycopodiella inundata (L.) Holub</i>	Northern bog club-moss	Lycopodiaceae	M	0	N	RB, vulnerable	-	Rare flavones chrysoseryl, segin and tricin; triterpenoid: o-onocerin; lysine-derived alkaloids; Lycodine type alkaloids, lycopodine alkaloids, tetracyclic alkaloids, clavine alkaloid, huperzine A; fatty oil, which consists of glycerides of higher fatty acids - palmitic, stearic, dioxystearic, oleic, linoleic and myristic, polymeric terpin, sporonin, phytosterol, glycerol, proteins, fiber, sugars and minerals; in the grass of plants plant are presence near 0.1-2% alkaloids, including lycopodine, clavatin, clavatoxine and lycodine. Antifungal, antibacterial, antiviral; relieve spasms, increase urine flow, estrogenic; reduce pain, fever, inflammation; insecticidal. Spores have anti-inflammatory, analgesic and diuretic effects and Used as a powder for infants, for the treatment of eczema, bedsores, wounds, burns and frostbite; At skin diseases - scabies, psoriasis, rashes, abscesses, boils; For local baths during seizures. In traditional Chinese medicine, an overground part of a plant that contains alkaloids is used to treat memory illnesses and Alzheimer's diseases.

<i>Lycopodium annotinum L.</i>	Stiff club-moss	Lycopodiaceae	M	0	N	RB, vulnerable	<p>Carotenoids, flavonoids, triterpenoids, nicotine; rare flavones chrysoeriol, selgin and tricin; triterpenoid: 21-episeratriol; lysine-derived alkaloids; lycodine type alkaloids, lycopodine alkaloids, tetracyclic alkaloids, clavine alkaloid, hyperizine A. Has hemostatic, weakening, diuretic, contraceptive, anticonvulsant, anesthetic, anti-inflammatory, astringent, insecticidal (veterinary) action.</p> <p>Antifungal, antibacterial, antiviral; relieve spasms, increase urine flow, estrogenic; reduce pain, fever, inflammation; insecticidal. Used for diseases of the stomach, liver, alcoholism, asthenia; for abrasions, eczema and hair loss.</p>
<i>Lycopodium clavatum L.</i>	Common club-moss	Lycopodiaceae	M	II	N	RP	<p>Fatty oil (spores), phenolic acids and their derivatives (spores), lycopoclavamine A and B, carotenoids, Flavonoids, sugars, minerals; rare flavones chrysoeriol, selgin and tricin; triterpenoid: o-onocerin; lysine-derived alkaloids; Lycodine type alkaloids, lycopodine alkaloids, tetracyclic alkaloids, clavine alkaloid, hyperizine A. Action - antitumor, sedative, antispasmodic, diuretic, contraceptive, anti-rheumatic, adsorbent, normalizes menstruation and the function of the gastrointestinal tract. Used in the treatment of metabolic, gout, rheumatic diseases, Alzheimer's disease, chronic kidney disease, mental illness.</p> <p>Antifungal, antibacterial, antiviral; relieve spasms, increase urine flow, estrogenic; reduce pain, fever, inflammation; insecticidal.</p>
						LC	

<i>Huperzia selago</i> (L.) Bernh. ex Schrank & C. Mart.	Northern fir-moss	Lycopodiaceae	R	0	F	RB, invaluable	LC	Selagoline, huperzine A and serratidine, carotene series β -C, Alpha-hoscurine, nankakurine A, acrifoline, des-N-methyl- β -hoscurine, huperzine, lycopodin, lycopodine, lycodine, cermizine B, lycoposerramine-G, lycopodine, fawcettidine, serratidine, lysine-derived alkaloids; Lycopodine type alkaloids, lycopodine alkaloids, tetracyclic alkaloids, clavine alkaloid. Antifungal, antibacterial, antiviral properties; relieve spasms, increase urine flow, estrogenic; reduce pain, fever, inflammation; insecticidal. Actions - vomiting, anticonvulsant, anti-inflammatory, diuretic, antitumor, analgesic. Used in the treatment of alcoholism, nicotine addiction, psoriasis, hysteria, neurasthenia, pulmonary tuberculosis, metabolic disorders.
<i>Selaginella</i> <i>helvetica</i> (L.) Spring.	Swiss spike- moss	Selaginellaceae	L	0	I	RB, extinct		Biflavonyl compounds (amentoflavone, sotetsuflavone); steroids, alkaloids, secolignans, neolignans, caffeine derivatives, alkaid glycosides, phenylpropanes, lignans. Used to treat diseases of the skin, gastritis, urinary tract, diabetes, hepatitis, cardiovascular problems
<i>Selaginella</i> <i>selaginoides</i> (L.) C. Mart.	Northern spike- moss	Selaginellaceae	R	II	F	RB, vulnerable	-	Biflavonyl compounds (amentoflavone, hinokiflavone, robustaflavone).
<i>Isoëtes lacustris</i> L.	Lake quillwort	Isoëtaceae	L	0	F	RB, vulnerable	-	Rare flavones chrysoeriol, selgin and tricin; luteolin bearing an extra hydroxyl group in 6-, 8- or 2'-position

After analyzing the State Register of Medicinal Products of Ukraine (14) we found out that most of the available drugs on the territory of Ukraine are manufactured and purchased abroad (Table 2).

In addition to the above-mentioned comprehensive homeopathic medications, there is also a

well-known cosmetic product - a powder on the basis of dry extract of *L. clavatum* (double-acting powder), which has a calming effect, and helps to reduce irritation and redness. The country of origin of the product is Israel.

*Table 2. Medications containing *Lycopodium* represented in the State Register of Medicines, Ukraine*

Name of the Drug	Active Substance	Producer	Applicant
METRO-ADNEX-INJEEL®	<i>Lycopodium clavatum</i>	Biologische Heilmittel Heel GmbH., Germany	Biologische Heilmittel Heel GmbH., Germany
NUX VOMICA-HOMACCORD®	<i>Lycopodium clavatum</i>	Biologische Heilmittel Heel GmbH., Germany	Biologische Heilmittel Heel GmbH., Germany
HEPEEL® N	<i>Lycopodium clavatum</i>	Biologische Heilmittel Heel GmbH., Germany	Biologische Heilmittel Heel GmbH., Germany
HEPEEL®	<i>Lycopodium clavatum</i>	Biologische Heilmittel Heel GmbH., Germany	Biologische Heilmittel Heel GmbH., Germany
PEKANA NATURHEILMITTEL.	<i>Lycopodium clavatum</i>	Biologische Heilmittel Heel GmbH., Germany	Biologische Heilmittel Heel GmbH., Germany
MOMORDICA COMPOSITUM	<i>Lycopodium clavatum</i>	Biologische Heilmittel Heel GmbH., Germany	Biologische Heilmittel Heel GmbH., Germany
TESTIS COMPOSITUM	<i>Lycopodium clavatum</i>	Biologische Heilmittel Heel GmbH., Germany	Biologische Heilmittel Heel GmbH., Germany
WEIGHT NORM	<i>Lycopodium clavatum</i>	PC "National Homeopathic Union", Ukraine	PC "National Homeopathic Union", Ukraine
HOLE-GRAN	<i>Lycopodium</i>	PC "National Homeopathic Union", Ukraine; Ferrer Internacional, S.A., Spain	PC "National Homeopathic Union", Ukraine; Ferrer Internacional, S.A., Spain
SCLERO-GRAN	<i>Lycopodium clavatum</i>	PC "National Homeopathic Union", Ukraine	PC "National Homeopathic Union", Ukraine
<u>GRAPHITES COSMOPLEX S</u>	<i>Lycopodium clavatum</i>	Biologische Heilmittel Heel GmbH., Germany	Biologische Heilmittel Heel GmbH., Germany
<u>ENERCEL® PLUS I. M</u>	<i>Lycopodium clavatum</i>	Laboratories Vijosa S.A. de C.V.	Laboratories Vijosa S.A. de C.V.
<u>ENERCEL® FORTE</u>	<i>Lycopodium clavatum</i>	Laboratories Vijosa S.A. de C.V.	Laboratories Vijosa S.A. de C.V.
<u>SOLVENCİUM</u>	<i>Lycopodium clavatum</i>	PC "Biolik"	LLC «Ukrainian pharmaceutical company»
<u>GEPATIUS "SPAG" - PEKANA</u>	<i>Lycopodium clavatum</i>	Pecana Heilmittel Heel GmbH., Germany	Pecana Heilmittel Heel GmbH., Germany

*Drugs are in bold and underlined - the term of registration has expired.

Medication containing Lycopods is also used in veterinary medicine. For example, Liarsin, which includes *L. clavatum*, is a complex homeopathic product, which has a corrective effect on protein, hydrocarbon, metabolism, restores impaired gastrointestinal function, stimulates the body's immune system. Its country of origin is Russia.

However, it should be noted that Lycopods have poisonous properties and their use should be carried out under medical supervision.

It is likely that this situation with the research and use of Lycopods is primarily conditioned by the absence of a raw material base in Ukraine and their conservation status.

CONCLUSION

The results obtained from the use of *Lycopodiophyta* indicate that there is no medical or any other economic importance in Ukraine. Most species of Lycopods are distributed only in certain regions and have a conservation status. The populations of the studied species are few and sensitive to anthropogenic factors. Most homeopathic preparations containing Lycopods are produced in other countries, which is caused by a number of factors listed above.

REFERENCES

1. Grodzinsky AM, editor. Medicinal plants. Kyiv: Encyclopedic Directory; 1992.
2. Kharchenko NS, Karamyshev AN, Sula VI, Voldarsky LI. Medicinal plants and their use. Kyiv: Zdorov'ia; 1981.
3. Lin CC, Kan WS. Medicinal plants used for the treatment of hepatitis in Taiwan. Am J Chinese Med. 1990; 18(1-2):35-43. doi: 10.1142/S0192415X9000006X.
4. Minarchenko VM. Medical vascular plants of Ukraine (medical and resource value). Kyiv: Fitosociocentre; 2005.
5. Nosal MA, Nosal IM. Medicinal plants and methods of their application in the people. Kyiv; 2013.
6. Grushvitsky IV, Zhilina SG, editors. Life of plants. V. 4. Mosses. Club mosses. Horsetails. Ferns. Gymnospermous plants. Moscow: Prosveshchenye; 1978.
7. Mosyakin SL, Fedoronchuk MM. Vascular plants of Ukraine: a nomenclatural checklist. Kiev; 1999.
8. Mosyakin SL, Tyshenko OV. A pragmatic phylogenetic classification of vascular Cryptogamic plants of the flora of Ukraine. Ukr Botan J. 2010; 67(6): 802-17.
9. Mukolaychuk VG. Botany. V. II. Course of lectures. Nikolaev: Nikolaev National Agrarian University; 2017.
10. Didukh YP, editor. Ecoflora of Ukraine. V. I. Kyiv: Fitodosociocentre; 2000.
11. Hassan HM, Jiang ZH, Syed TA, Qin W. Northern Ontario medicinal plants. Can J Plant Sci. 2012; 92(5):815-28. doi:10.4141/cjps2012-006.
12. Prokypin JD, editor. The determinant of higher plants of Ukraine. Kyiv: Naukova dymka; 1987.
13. Didukh YP, editor. Red Book of Ukraine. Kyiv: Globalkonsaltyng; 2009.
14. State Register of Medicinal Products of Ukraine, Available at: <http://www.drlz.com.ua/>
15. Minarchenko VM, Tymchenko IA. Atlas of medicinal plants of Ukraine Kyiv: Fitodosociocentre; 2002.
16. Tsvelev NN, editor. Conspectus Flora Europae Orientalis. Tomus 1. Petropoli-Mosqua: Consociatio editionum scientificarum KMK; 2012.
17. Allen D, Bilz M, Leaman DJ, Miller RM, Timoshyna A, Window J. European Red List of Medicinal Plants. Luxembourg: Publications Office of the European Union; 2014.
18. Bilz M, Kell SP, Maxted N, Lansdown RV. European Red List of Vascular Plants. Luxembourg: Publications Office of the European Union; 2011.
19. IUCN. Red List categories and criteria: Version 3.1. 2nd ed. Gland (Switzerland); Cambridge (UK): IUCN; 2012.
20. Ayer W. The Lycopodium alkaloids. Nat Prod Rep. 1991; 8(5):455-63. doi: 10.1039/np9910800455.
21. Czapski GA, Szypuła W, Kudlik M, Wileńska B, Kania M, Witold Danikiewicz W, et al. Assessment of antioxidative activity of alkaloids from *Hyperzia selago* and *Diphasiastrum complanatum* using in vitro systems. Folia Neuropathol. 2014; 52(4):394-406. doi: 10.5114/fn.2014.47840.
22. Czeczuga B. Carotenoids in sixty-six representatives of the Pteridophyta. Biochem Syst Ecol. 1985b; 13(3):221-30. doi:10.1016/0305-1978(85)90030-4.
23. Kataoka K, Mito H, Kogure N, Kitajima M, Wongseripatana S, Arisawa M, et al. Ten new fawcettimine-related alkaloids from three species

- of *Lycopodium*. *Tetrahedron*. 2011; 67(35):6561-7. doi:10.1016/j.tet.2011.05.107.
24. Ma X, Gang DR. The *Lycopodium* alkaloids. *Nat Prod Rep*. 2004; 21(6):752-72. doi: 10.1039/b409720n.
25. Filho JMB, Alencar AA, Nunes XP, de Andrade Tomaz AC, Sena-Filho JG, Athayde-Filho PF, et al. Sources of alpha-, beta-, gamma-, delta- and epsilon-carotenes: A twentieth century review. *Braz J Pharmacogn*. 2008; 18(1):135-54. doi:10.1590/S0102-695X2008000100023.
26. Flora of North America Editorial Committee, editors. *Flora of North America North of Mexico*. Vol.2. New York & Oxford: Oxford Univ Press; 1993. Available at: http://www.efloras.org/volume_page.aspx?volume_id=1002&flora_id=1
27. Halldorsdottir ES, Palmadottir RH, Nyberg NT, Olafsdottir ES. Phytochemical analysis of alkaloids from the Icelandic club moss *Diphasiastrum alpinum* (L.) Holub. *Planta Med*. 2012; 78(11):402. doi: 10.1055/s-0032-1321089.
28. Ishiuchi K, Kubota T, Ishiyama H, Hayashi S, Shiba T, Mori K, et al. Lyconadins D and E, and complanadine E, new *Lycopodium* alkaloids from *Lycopodium complanatum*. *Bioorg Med Chem*. 2011;19(2):749-53. doi: 10.1016/j.bmc.2010.12.025.
29. Ma X, Tan Ch, Zhu D, Gang DR, Xiao P. Huperzine A from *Huperzia* species - An ethnopharmacological review. *J Ethnopharmacol*. 2007; 113(1):15-34. doi: 10.1016/j.jep.2007.05.030.
30. Oirin B, Jay M. Etude chimiosystematique des Lycopodiales, Isoetales Selaginellales et Psilotales. *Biochem Syst Ecol*. 1978b; 6(2):99-102. doi:10.1016/0305-1978(78)90032-7.
31. Harborne JB, Baxter H, Moss GP, editors. *Phytochemical Dictionary. A Handbook of Bioactive Compounds from Plants*. 2nd Edition. London; Philadelphia: Taylor&Francis; 1999.
32. Sokolov VS. Alkaloidonic plants of the USSR. AN USSR; 1952.
33. Voirin B. Distribution des composés polyphénoliques chez les Lycopodiées. *Phytochemistry*. 1972; 11(1):257-62. doi:10.1016/S0031-9422(00)90000-1.
34. Voirin BJM, Hauteville M. Isoetine, nouvelle flavone isolée de Isoetes delilei et de Isoetes du-rieui. *Phytochemistry*. 1975; 14(1):257-60. doi:10.1016/0031-9422(75)85050-3.
35. Voirin BJM, Hauteville M. Selagine, nouvelle flavone isolée de *Huperzia selago*. *Phytochemistry*. 1976; 15:840-1.
36. Voirin BJM. Apport de la Biochimie Flavonique à la systematique du genre *Lycopodium*. *Biochem Syst Ecol*. 1978a; 6(2):95-7. doi:10.1016/0305-1978(78)90031-5.
37. Voirin B. Recherches chimiques, taxinomiques et physiologiques sur les flavonoïdes des Pteridophytes. Thèse. Docteur-Science. L'Université de Lyon; 1970.
38. Tsuda Y, Fujimoto T, Isobe K, Sano T, Kobayashi M. Chemotaxonomical studies on the triterpenoids of *Lycopodium* plants. *Yakugaku Zasshi*. 1974; 94(8):970-90.